

EXHIBIT F

REVIEWS

Safety considerations for synthetic sling surgery

Jerry G. Blaivas, Rajveer S. Purohit, Matthew S. Benedon, Gabriel Mekel, Michael Stern, Mubashir Billah, Kola Olugbade, Robert Bendavid and Vladimir Iakovlev

Abstract | Implantation of a synthetic midurethral sling (SMUS) is the most commonly performed anti-incontinence operation in women worldwide. The effectiveness of the SMUS is comparable to that of the historical gold standards—autologous fascial slings and the Burch colposuspension. Much controversy, however, has evolved regarding the safety of this type of sling. Overall, the quality of the studies with respect to assessing risks of SMUS-associated complications is currently poor. The most common risks in patients with SMUS include urethral obstruction requiring surgery (2.3% of patients with SMUS), vaginal, bladder and/or urethral erosion requiring surgery (1.8%) and refractory chronic pain (4.1%); these data likely represent the minimum risks. In addition, the failure rate of SMUS implantation surgery is probably at least 5% in patients with stress urinary incontinence (SUI). Furthermore, at least one-third of patients undergoing sling excision surgery develop recurrent SUI. Considering the additional risks of refractory overactive bladder, fistulas and bowel perforations, among others, the overall risk of a negative outcome after SMUS implantation surgery is $\geq 15\%$.

Blaivas, J. G. *et al.* *Nat. Rev. Urol.* advance online publication 18 August 2015; doi:10.1038/nrurol.2015.183

Introduction

The Burch colposuspension and autologous fascial pubovaginal sling have been considered the gold standard treatments of stress (sphincteric) incontinence (SUI), in women since the late 1990s.¹ Historically, pubovaginal slings had been reserved for women with complicated, severe and/or recurrent sphincteric incontinence,² but since the late 1990s these have been advocated for treatment of women with all types of sphincteric incontinence—simple and complicated.^{3,4} Over the past decade, fueled by a stampede of innovations in synthetic sling composition, structure and implantation techniques and a surge in commercial marketing, implantation of the synthetic midurethral sling (SMUS) has emerged as the most frequently performed operation in women with SUI. Some authors suggest that, to date, over 3 million SMUS implantation procedures have been conducted worldwide, and more than 80% of these happened in the USA.⁵ We have been unable to independently verify the number of SMUS implanted worldwide. But, by extrapolating the data from a population-based cohort study,⁶ we estimate that approximately 500,000 SMUS were implanted in the USA between years 2001–2010 and that in the ensuing 4 years at least another 300,000 of these procedures were done. Considering the size of the population of the rest of the world and the fact that slings have been implanted *en masse* in most economically developed countries since at least the mid-1990s, a figure of 3 million procedures seems reasonable.⁶

Furthermore, in an analysis of 7,200 case logs submitted by American urologists for their certifying credentials in 2013, 83% of operations performed for incontinence in women were SMUS implantations.⁷

SMUS implantation is an operation for the correction of sphincteric incontinence in which a synthetic plastic like mesh strip (the sling) is passed around the urethra into the retropubic space or beneath the urethra through the obturator foramen using trocars. Theoretically, when abdominal pressure rises, as in a cough or sneeze, the urethra is compressed by the sling, and the flow of urine is prevented, much like kinking of a garden hose. The appeal of such procedures is obvious—in theory. SMUS implantation is a minimally invasive, easy-to-perform procedure that is usually completed in under half an hour and, compared to traditional native tissue repairs, enables a much faster recovery with less perioperative morbidity than either the Burch colposuspension or autologous fascial slings. The effectiveness of this approach remains unchallenged. Numerous trials have shown SMUS to be as effective as the autologous fascial sling and Burch colposuspension, with moderate and/or high quality of evidence.⁸

Theory and practice often diverge, though, and this seems to be the case with SMUS. As more SMUS implantations are being performed and the longevity expectations of patients with SMUS are increasing, it has become apparent that unanticipated, serious, sometimes lifestyle-altering complications can occur that are not only unique to patients with SMUS but are also often refractory to treatment.^{9,10} The purpose of this Review is to summarize the published literature regarding complications that are uniquely associated with SMUS and to present an overview of complications that are not unique to these slings.

Competing interests

J.G.B. and V.I. have provided opinions as medicolegal expert witnesses in mesh litigation cases. J.G.B. has acted as a consultant for Astellas Pharma, is a shareholder in P Square Medical, and is a shareholder and co-owns intellectual property with LLC and Sympelligence Medical Informatics. The other authors declare no competing interests.

Department of Urology (J.G.B., R.S.P.), Institute for Bladder and Prostate Research (M.S.B.), Weill Cornell Medical College, Cornell University, 445 East 77th Street, New York, NY 10075, USA. Hahnemann University Hospital, 230 North Broad Street, Philadelphia, PA 19102, USA (G.M.). SUNY Downstate College of Medicine (M.S., M.B.), SUNY Downstate Medical Center (K.O.), 450 Clarkson Avenue, Brooklyn, NY 11203, USA. Shouldice Hospital, 7750 Bayview Avenue, Thornhill, ON L3T 4A3, Canada (R.B.). St Michael's Hospital, University of Toronto, 30 Bond Street, ON M5B 1W8, Canada (V.I.).

Correspondence to: J.G.B. jgblys@gmail.com

Key points

- The effectiveness of synthetic midurethral slings (SMUS) is comparable to the time-honoured gold standards—the autologous fascial sling and Burch colposuspension
- At least 15% of women with SMUS experience a serious adverse outcome and/or recurrent sphincteric incontinence
- A subset of women sustain refractory, lifestyle-altering complications that are unique to women with a SMUS
- SMUS-associated complications are under-reported
- The overall quality of the published evidence is currently low with respect to assessing SMUS safety and SMUS-associated complications

Transvaginal mesh slings

The retropubic tension-free transvaginal mesh tape (RP) sling procedure was introduced for treatment of SUI in 1995,¹¹ and was soon followed in 2001 by the transobturator tape (TOT) sling procedure, in which the sling is introduced via the obturator fossa instead of the retropubic space in an attempt to minimize the complications associated with use of RP slings.¹² Since the introduction of these procedures, results of many modifications of the RP and TOT sling procedures have been published, generally with a follow-up duration of <4 years; most studies had a follow-up duration of ≤1 year.

In a review of prospective randomized controlled trials investigating performance of RP and/or TOT slings, authors reported objective cure rates of patients treated with RP slings ranging between 83.9% and 100%, and 84% and 97.6% for those with TOT slings. Subjective cure rates for patients with RP or TOT slings were 64.5–94% and 60–92.9%, respectively. Median or mean follow-up duration of these studies ranged between 9 months and 24 months.¹³ Thus, no notable differences in effectiveness have been revealed by meta-analyses of trials comparing the effectiveness of RP slings with that of TOT slings.^{14,15}

More than 19 years have elapsed since the introduction of RP slings and 13 years for TOT slings; results regarding the long-term effectiveness of these treatments are, therefore, now available, with the longest series reporting a follow-up duration of 17 years.¹⁶ Findings of over 200 studies of varying design and quality, investigating the effectiveness of either RP or TOT slings have been published. Of these, only six contain information on outcomes of patients followed up for >5 years and five have reported outcomes after a follow-up duration of 5 years (Table 1). Effectiveness of the treatments tested in these studies has been measured using a variety of subjective and/or objective outcome instruments. Subjective outcomes have been measured using detailed validated questionnaires such as the short urogenital distress inventory (SUDI), short incontinence impact questionnaire, the European quality of life questionnaire,¹⁷ the patient global impression of improvement (PGII) questionnaire,^{16,18} the visual/verbal analogue scale (VAS),^{16,19–21} non-validated questionnaires^{20,22} and telephone surveys.²³ In our judgement, however, the use of some of the validated outcome instruments does not provide sufficiently explicit data

to be considered scientifically rigorous. For example, the urogenital distress inventory (UDI)-6 conflates bother with incontinence. This inventory contains the question “Do you experience and how much are you bothered by ... leakage,”²⁴ from stress incontinence? This means a woman could leak only a few drops once a year and have a lot of bother or be totally incontinent and have no bother, yet record exactly the same score. Furthermore, variability in the use of validated and nonvalidated questionnaires might be another explanation of the discrepancies in reported outcomes and can complicate direct comparisons of results from different studies.

A variety of objective outcome instruments have also been used to measure outcomes of these long-term (follow-up duration ≥5 years) studies. The Valsalva stress test, in which the patient is asked to perform a Valsalva manoeuvre to increase abdominal pressure, or the cough test with a full bladder to provocatively test for the development of SUI are the most commonly used objective outcome measures.^{17,18,20,21} However, these tests are not standardized and, typically, abdominal and vesical leak point pressures are not measured.²⁵ Other studies involved checking bladder volume by ultrasonography or just verbally confirming that the patient feels that their bladder is “comfortably full”²⁶ or “full”^{18,27} and then asking the patient to cough. One group asked the patient to do “20 jumping jacks and 3 forceful coughs”²² or “10 coughs in the standing position”²² with a 300 ml bladder volume to check for SUI.²²

The pad-weight test is another objective measurement used to assess treatment outcomes. A variety of pad-weight tests were used in these long-term studies including a 1-hour pad-weight test,¹⁹ a 24-hour pad-weight test²⁸ and a 48-hour pad-weight test²⁹ with a result of ≤1 g increase in pad weight defining cure. Two studies used urodynamic evidence of the absence of leak on performing a Valsalva manoeuvre to indicate post-operative success either as a primary endpoint or as an adjunctive measure.^{17,18}

Of published series with follow-up durations ≥5 years, the largest consisted of 707 patients and the smallest consisted of 55 patients.^{16–23,26,27,29} Overall, objective cure rates of patients with SUI after treatment with either TOT or RP slings at or after a follow-up duration of 5 years ranged between 71% and 92%, and subjective cure rates between 65% and 90.3%. By combining subjective measures of cure and improvement, treatment effectiveness increased from 76% to 95% (Table 1).^{16–23,26,27,29}

The series with the longest reported follow-up duration (17 years) was a prospective, single-institution study of 90 women who underwent the (original) RP sling implantation procedures at Uppsala University.¹⁶ The overall rate of subjective cure or improvement assessed using the PGII score was 87% and the objective cure rate assessed using a cough stress test was 91%. Between follow-up years 5 and 17, objective cure rates declined very little—from 94% to 91%—and subjective cure or improvement decreased from 95% to 87%. However, 11 of 53 evaluable patients said that they

Table 1 | Long-term (follow-up duration >5 years) studies of SMUS effectiveness

Study characteristics	Patient characteristics	Mean follow-up duration (months)	Outcome Instrument	Outcomes*
Prospective studies				
Angioli <i>et al.</i> (2010) ²⁰ <i>n</i> = 72	Outcomes of 69 patients with RP or TOT slings were evaluated; 4.1% were lost to follow up	60	ST, NVQ, [‡] VAS [‡]	Objective cure reported in 71% and 73% in patients with RP or TOT slings, respectively; Subjective cure reported by 60% and 62% of patients with RP or TOT slings, respectively
Groutz <i>et al.</i> (2011) ²³ <i>n</i> = 60	Outcomes of 52 patients with RP slings were evaluated; 13.3% were lost to follow up	60	NVQ [‡]	Subjective cure reported by 65% of patients
Groutz <i>et al.</i> (2011) ²⁷ <i>n</i> = 65	Outcomes of 61 patients with TOT slings were evaluated; 6.1% were lost to follow up	60	ST, NVQ [‡]	Objective cure reported in 74% of patients, 8% had improved symptoms and 18% subjectively reported treatment failure
Cheng <i>et al.</i> (2012) ¹⁷ <i>n</i> = 103	Outcomes of 100 patients with TOT slings were evaluated; 2.9% were lost to follow up	65	VUD, QOL, [‡] VAS [‡]	Objective cure reported in 87.4% of patients; subjective cure reported by 78% of patients
Nilsson <i>et al.</i> (2013) ¹⁶ <i>n</i> = 90	Outcomes of 58 patients with RP slings were evaluated; 23.3% were lost to follow up	201	ST, VQ [‡]	Objective cure reported in 91.3% of patients; subjective cure reported by 87%
Serati <i>et al.</i> (2013) ¹⁸ <i>n</i> = 191	Outcomes of 185 patients with TOT slings were evaluated; 3.1% were lost to follow up	60	ST, VQ [‡]	Objective cure reported in 90.3% of patients; subjective cure reported by 90.8%
Svenningsen <i>et al.</i> (2013) ²² <i>n</i> = 603	Outcomes of 483 patients with RP slings were evaluated; 19.9% were lost to follow up	120	Exercise + PT, VQ, [‡] NVQ [‡]	Objective cure reported in 89.9% of patients; subjective cure reported in 76.1% of patients; 18% had improved symptoms; 5.9% had treatment failure
Retrospective studies				
Ankardal <i>et al.</i> (2006) ²⁹ <i>n</i> = 707	Outcomes of 271 patients with RP slings were evaluated; 5.0% were lost to follow up	60 [§]	ST, 48 h PT (NVQ, [‡] VAS [‡])	Objective cure reported in 83% of patients; subjective cure reported by 73.1% of patients; 15.9% had improved symptoms; 11% had treatment failure
Olsson <i>et al.</i> (2010) ²¹ <i>n</i> = 147	Outcomes of 104 patients with RP slings were evaluated; 15.6% were lost to follow up	138	ST	Objective cure reported in 84% of patients; subjective cure reported by 77% of patients; 18% had improved symptoms; 5% had treatment failure
Li <i>et al.</i> (2012) ¹⁹ <i>n</i> = 55	Outcomes of patients with RP slings were evaluated; percentage of patients lost to follow up not reported	81.85	1 h PT (NVQ [‡])	Objective cure reported in 85.5% of patients; subjective cure reported in 74.6% of patients; 7% had improved symptoms; 25.6% had treatment failure
Athanasίου <i>et al.</i> (2014) ²⁶ <i>n</i> = 145	Outcomes of 124 patients with TOT slings were evaluated; 14.4% were lost to follow up	90.3	ST (VQ [‡])	Objective cure reported in 81.5% of patients; subjective cure reported in 83.1% of patients; 3.2% had improved symptoms; 13.7% had treatment failure

*Owing to a lack of uniformity in reporting efficacy (improved and failed), improvement and failure were assumed to be based on subjective responses. Incidence of failure was calculated by subtracting the sum of subjective cured and improved responses from 100%. Improved patients were mutually exclusive to cured patients reported. †Indicates a subjective outcome instrument. §Indicates actual, not mean follow-up duration. Abbreviations: NVQ, nonvalidated questionnaire; PT, pad-weight test; QOL, quality of life; RP, retropubic tension-free transvaginal mesh tape; SMUS, synthetic midurethral slings; ST, cough or valsalva stress test; TOT, transobturator tape; VAS, visual analogue scale; VQ, validated questionnaire; VUD, videourodynamics.

“experienced leakage during straining”.¹⁶ The authors attributed this symptom to the development of severe urge incontinence, which might or might not have been caused by the RP sling itself. However, this study was hindered by the fact that 32 of 90 patients (36%) were not included in the analysis after a follow-up duration of 17 years as 11 (12%) women died, five (6%) had mental impairment and 16 (18%) were lost to follow-up. Overall, 58 (64%) women were available to have their 17-year outcomes evaluated, of whom 46 (51%) women were evaluated in the clinic and 12 (13%) were interviewed by telephone.

By contrast, in another investigation,³⁰ researchers using much more stringent outcome criteria found the 2-year objective success rate of RP and TOT slings was 77% and 72.3%, respectively and the subjective success rates were 56% and 48%—nearly a 40% reduction in subjective success compared with results from the Uppsala cohort.¹⁶ In this randomized study of 597 patients, objective success was defined as a negative provocative stress test, a negative 24-hour pad-weight test

and no need for retreatment of SUI.²⁷ Subjective success was defined as the absence of self-reported symptoms of SUI on the Medical, Epidemiological and Social Aspects of Aging questionnaire and no urine leakage recorded in a 3-day voiding diary.

The overall effectiveness of RP or TOT slings reported in these long-term studies of patients with SUI has been generally high, although some of the reported success occurred after a secondary operative procedure or medical intervention.^{17,19,21,26} Thus, for patients who developed mesh erosion and had successful revision surgery, studies would typically report this as a successful outcome. Secondary treatments for those who develop *de novo* urge incontinence have not been reported, although if the RP or TOT sling procedures are effective in these patients they would typically be included in the ‘subjectively cured’ or ‘improved’ category.

The percentage of patients who were lost to follow up should also be carefully noted. In patients who were lost to follow up because of death, the cause of death and

Table 2 | Complications of either RP or TOT slings

Complication	<i>n</i>	Complications (% of patients)	Incidence (mean; range)
General complications			
Death within 30 days*	7,762	0 (0.0)	0.0
Urethral obstruction/voiding dysfunction	25,586	1,403 (5.5)	7.3; 0–33.9
Urethral obstruction requiring surgery	9,375	301 (3.2)	2.3; 0–21.3
Urinary infections	13,296	598 (4.5)	7.3; 0–39.1
Pain (within 6 weeks)	5,097	374 (7.3)	7.9; 0–33.3
Neurologic symptoms (within 6 weeks)	1,769	42 (2.4)	1.2; 0–10.3
<i>De novo</i> OAB	14,765	1,512 (10.2)	10.9; 0–48.1
Pelvic organ perforation			
In total	20,630	681 (3.3)	3.5; 0–16.1
Bladder	19,411	579 (3.0)	2.9; 0–16.1
Vaginal	5,521	91 (1.6)	1.4; 0–14.1
Urethral	4,541	6 (0.1)	0.0; 0–1.5
Bowel	3,820	4 (0.1)	0.0; 0–1.7
Ureteral	3,820	1 (0.0)	0.0; 0–1.3
Mesh exposure/erosion/extrusion			
In total	17,520	475 (2.7)	2.5; 0–26.1
Treated conservatively	15,403	112 (0.7)	0.9; 0–7.1
Vaginal	13,496	78 (0.6)	0.7; 0–7.1
Bladder	13,496	5 (0.0)	0.0; 0–5.6
Urethral	13,496	0 (0.0)	0.0
Requiring surgery	16,619	333 (2.0)	1.8; 0–26.1
Vaginal	13,705	235 (1.7)	1.5; 0–15.9
Bladder	13,393	29 (0.2)*	0.2; 0–15.2
Urethral	13,628	11 (0.1)	0.2; 0–16.7
Longer-term complications			
Refractory pain (>6 weeks)	7,084	247 (3.5)	4.1; 0–30.5
Neurologic Symptoms (>6 weeks)	2,449	51 (2.0)	1.0; 0–10.6
Fistulas	710	2 (0.3)	0.3; 0–1.1

*No deaths were reported in peer-reviewed publications, although 7 were reported in the MAUDE database.
*Three studies removed from incidence calculation because these were case series of just bladder erosions. Abbreviations: MAUDE, manufacturer and user facility device experience; OAB, overactive bladder; RP, retropubic tension-free transvaginal mesh tape; TOT, transobturator tape.

whether it might have been related to having a TOT or RP sling has not been reported. In addition, long-term studies often featured a lack of follow up in a substantial number of patients. In these long-term studies, between 5% and 36% of patients were either deceased or unavailable for follow up for other reasons (Table 1).

In summary, the long-term effectiveness of RP or TOT slings, as measured by subjective and/or objective instruments suggests that rates of cure or improvement of SUI after implantation are high and compare favourably to the traditional gold standard—the autologous pubovaginal sling. These results might, however, be overly optimistic owing to a host of factors including the suboptimal outcome instruments used, inclusion of patients who might have required multiple procedures and the loss of a substantial number of patients to follow-up monitoring.

SMUS complications

Mesh sling complications can be caused by a host of factors: intraoperative transgressions (such as viscus or vaginal perforation and nerve injury);³¹ bacterial contamination;³² improper tensioning of the sling—either too tight or too loose; host–foreign-body reaction;³³ tissue ingrowth;^{34,35} and changes that the mesh undergoes once implanted, such as degradation, curling, banding and leaching of substances into the surrounding tissues (Tables 2–5).^{36–38}

Evaluating the incidence, severity and consequences of the various RP and/or TOT sling complications is a daunting task. Of the thousands of published studies, only a few were even designed to track complications in any meaningful way. The short follow-up duration of most of these studies and the lack of accounting for those lost to follow up are additional confounders. In addition, complications might arise that were not even recognized when the original studies were conducted, such as banding as a cause of dyspareunia, which was first described in 2010.³⁸ Furthermore, all studies are hampered by an absence of the patient's own perception of the severity of the complication. For example, one study³⁹ that included pain lasting >6 weeks as a category of complication was published, but there is no mention of the severity of this pain, effect on quality of life nor how long the pain actually lasted. For some patients, this pain is treatment-refractory and lifestyle-altering, yet no metric exists that describes this category of complication in sufficient detail. The effects of long-term pain receive no attention at all in any study except for a few case studies of complications.^{9,40–44}

The concomitant, widespread use of two different generic sling designs (RP and TOT) with different implantation techniques and at least 41 different commercially available kits,⁴² each having different sling and trocar characteristics with potentially different complication profiles, confounds accurate analysis of sling complications. These different characteristics might also portend different complication profiles, yet studies of sling complications almost never distinguish between the different kit types and many do not even separate TOT from RP slings.

Considerable evidence exists that SMUS complications are underreported. Discrepancies exist between the SMUS complication rates reported by urologists from individual institutions, those reported in the literature, the (unreported) experience of tertiary care practices and those in the MAUDE (Manufacturer and User Facility Device Experience) database.^{10,40,45} For example, in a population-based cohort of 45 million commercially insured individuals in the USA, investigators found the cumulative risk of requiring sling removal owing to voiding dysfunction or mesh extrusion and/or erosion to be 3.7% (95% CI 3.5–3.9%) after a follow-up duration of 9 years.⁶ Furthermore, this study⁶ excluded patients whose slings were removed owing to pain and other indications, thus the actual incidence of sling removal owing to complications is probably even higher than that. Extrapolating from this estimate and the estimated number of slings implanted in the

Table 3 Complications of RP slings			
Complication	n	Complications (% of patients)	Incidence (mean; range)
General complications			
Death within 30 days	3,499	0 (0.0)	0.0
Urethral obstruction/voiding dysfunction	16,301	704 (2.8)	8.8; 0–32.7
Requiring surgery	6,875	223 (2.4)	2.7; 0–8.9
Urinary infections	8,936	327 (3.7)	8.6; 0–39.1
Pain (within 6 weeks)	2,133	111 (5.2)	4.5; 0–23.1
Neurologic symptoms (within 6 weeks)	520	14 (2.7)	1.6; 0–5.0
<i>De novo</i> OAB	7,989	925 (11.6)	11.4; 0–29.4
Pelvic organ perforation			
In total	13,164	498 (3.8)	4.8; 0–14.3
Bladder	12,929	480 (3.7)	4.6; 0–14.3
Vaginal	763	11 (1.4)	1.0; 0–2.1
Urethral	1,224	4 (0.3)	0.0; 0–1.5
Bowel	800	4 (0.5)	0.0; 0–1.7
Ureteral	800	0 (0.0)	0.0
Mesh exposure/erosion/extrusion			
In total	8,303	179 (2.2)	2.3; 0–26.1
Treated conservatively	7,168	44 (0.6)	0.1; 0–5.6
Vaginal	6,193	22 (0.4)	0.0; 0–4.6
Bladder	6,193	5 (0.1)	0.0; 0–5.6
Urethral	6,193	0 (0.0)	0.0
Requiring surgery	7,902	135 (1.7)	1.6; 0–26.1
Vaginal	6,621	79 (1.2)	1.0; 0–10.9
Bladder	6,386	26 (0.4)	1.4; 0–15.2
Urethral	6,621	4 (0.1)	0.3; 0–16.7
Longer-term complications			
Refractory pain (>6 weeks)	2,328	42 (1.8)	2.0; 0–7.9
Neurologic symptoms (>6 weeks)	908	19 (2.1)	1.0; 0–5.2
Fistulas	388	1 (0.2)	0.4; 0–0.7
Abbreviations: OAB, overactive bladder; RP, retropubic tension-free transvaginal mesh tape.			

USA (80% of 3 million slings worldwide),⁵ approximately 88,000 removal procedures should have occurred, yet we could find only 740 such procedures that have been reported in peer-reviewed publications.^{17,18,20,26,31,43,46–99} An additional 7,654 mesh removals are reported in patient series investigating sling complications.^{6,9,10,41,44,100–114} Use of imperfect research methodologies, a lack of long-term follow up and reporting bias have been suggested as causes of these differences.^{10,45,115,116}

Safety and risk:benefit considerations

Safety of SMUS surgery refers to the probability of any adverse event, while risk describes the range and probability of specific adverse events. Demonstrating risk is relatively easy, but assessing safety is much more difficult. Any known adverse event is a risk, regardless of how infrequently the event has been reported or observed. Even a single case report of an adverse event establishes the existence of a specific risk, although without knowing

the denominator, accurate assessments of the safety of sling surgery are impossible. The problem, simply stated, is that no well-controlled long-term safety studies with published results currently exist, nor do any good registries in this area. In lieu of this lack of conclusive evidence, we present current data, which, at its best, represent the minimum risks associated with SMUS implantation and long-term use (Tables 1 and 2). Major risks of SMUS surgery, which should be weighed up by patients considering undergoing these procedures, include those requiring further surgery and those that are refractory to treatment. Complications that lead to further surgery include urethral obstruction (3.2%), vaginal, bladder and/or urethral erosion (2%) and fistulas (0.3%). In addition, we estimate that bowel perforation and serious infections have a combined incidence of about 0.1%.^{117–134} Refractory and potentially lifestyle-altering complications include chronic pain (4.1%) and *de novo* overactive bladder (OAB) in 11% (Table 5), although the number of patients with *de novo* OAB who are refractory to treatment remains unknown. Evidence suggests that well over 50% of patients with OAB of any aetiology discontinue medical treatment within 1 year, owing to a combination of poor effectiveness and low tolerability.^{135–137} For the purposes of this discussion we made a conservative estimate that 35% of patients with *de novo* OAB following SMUS surgery are refractory to treatment, which suggests that approximately 3.9% of patients who have undergone SMUS surgery will have refractory OAB (Box 1).

Establishing safety, defined as the chances of having an unsatisfactory outcome following sling surgery, is an important step in enabling accurate patient decision making. Many patients with both chronic pain and *de novo* OAB have previously undergone SMUS revision surgery; however, simply adding together the complication incidences to give an accurate indication of safety is impossible, as the available data are not sufficiently reliable to produce an accurate estimate this way. For the purposes of this discussion, we have made our best estimate of the safety of SMUS surgery utilizing the data summarized in the preceding paragraph.

The reported 5-year failure rate of SMUS surgery in patients with SUI ranges from 5% to 26% (Table 1),^{16–23,26,27,29} and the reported incidence of *de novo* SUI after sling excision surgery ranges from 10–62%.^{9,43,78,84,85,94,100,138} From these data, we estimate the lowest rate of recurrent and/or persistent SUI among patients who underwent SMUS surgery to be 5.3%. Furthermore, we conclude that the lowest estimated risk of serious complications of SMUS surgery is 13.6% and the additional risk of failure with respect to the original procedure (with respect to SUI) is 5.3%. These estimated data reflect the minimum risks reported in the literature, the actual risks could be considerably higher.

The ability of both physicians and patients to make informed decisions regarding sling surgery is predicated on an accurate understanding of the risk:benefit ratio. The benefits of sling surgery, with respect to effectiveness, have been well documented. By contrast, the risks associated with SMUS surgery are poorly understood

Table 4 | Complications of patients with a TOT sling

Complication	n	Complications (% of patients)	Incidence (mean; range)
General complications			
Death within 30 days*	4,044	0 (0.0)	0.0
Urethral obstruction/voiding dysfunction	8,287	406 (4.9)	5.9; 0–33.9
Requiring surgery	5,001	75 (1.5)	2.0; 0–21.3
Urinary infections	4,003	226 (5.6)	6.2; 0–23.3
Pain (within 6 weeks)	2,964	262 (8.8)	10.2; 0–33.3
Neurologic symptoms (within 6 weeks)	1,249	28 (2.2)	0.9; 0–10.3
De novo OAB	6,215	519 (8.4)	10.3; 0–48.1
Pelvic organ perforation			
In total	5,856	143 (2.4)	2.3; 0–16.1
Bladder	4,872	60 (1.2)	1.1; 0–16.1
Vaginal	4,582	80 (1.7)	1.6; 0–14.1
Urethral	4,296	2 (0.0)	0.0; 0–0.3
Bowel	3,020	0 (0.0)	0.0
Ureteral	3,020	1 (0.0)	0.0; 0–1.3
Mesh exposure/erosion/extrusion			
In total	8,293	196 (2.4)	2.7; 0–19.0
Treated conservatively	7,648	64 (0.8)	1.0; 0–7.1
Vaginal	6,739	52 (0.8)	1.0; 0–7.1
Bladder	6,739	0 (0.0)	0.0
Urethral	6,739	0 (0.0)	0.0
Requiring surgery	7,901	132 (1.7)	1.9; 0–15.9
Vaginal	6,528	98 (1.5)	1.8; 0–15.9
Bladder	6,528	3 (0.0)*	0.0
Urethral	6,528	7 (0.1)	0.0; 0–2.6
Longer-term complications			
Refractory pain (>6 weeks)	4,756	204 (4.3)	5.3; 0–30.5
Neurologic symptoms (>6 weeks)	1,541	32 (2.1)	0.9; 0–10.6
Fistulas	322	1 (0.3)	0.2; 0–1.1

*Case series only investigating bladder erosions were removed from the analysis. Abbreviations: OAB, overactive bladder; TOT, transobturator tape.

and poorly documented. Some risk factors, such as prior pelvic radiation or surgery, concomitant urethral diverticular surgery, the surgical learning curve and an individual surgeon's skill set are accepted risk factors, yet even these risk factors are not well documented in peer-reviewed publications. Thus, accurately prognosticating the risks of SMUS surgery for any particular patient is difficult.

Owing to the limitations inherent in the evaluation of risk and safety described in this section, we can estimate that, based on the available literature reports, a minimum of 12.5% of women who undergo mesh SMUS surgery have a serious adverse event and/or surgical failure, although limited data are available on prognostic indicators.

Perioperative complications

Methods used to identify and classify perioperative complications vary widely between studies. A classification

system originally devised in 2004 based on severity of complications¹³⁹ was modified for use by the Urinary Incontinence Treatment Network, a multicentre collaboration supported by the National Institute of Diabetes and Digestive and Kidney Diseases, a branch of the NIH.^{30,39,140,141} Investigators in this network defined minor complications (or minor adverse events) as those not requiring surgical intervention that were treated expectantly or with medication (grade 1–2). Major complications (serious adverse events) include those requiring one or more surgical procedures, and life-threatening complications are defined as those requiring management in an intensive care unit and often resulting in patient death (grade 3–5).³⁰ Bladder or urethral trocar perforation was considered to be a serious adverse event whether or not further intervention was necessary. Another classification scheme, the Accordion system,¹²⁰ enables postoperative complications to be categorized into in four levels of severity: mild, moderate, severe, and death.^{44,142}

However, sorting complications into groups based solely on the severity of their presentation and treatment might be misleading in the absence of adequate follow-up monitoring. For example, patients with apparently minor complications such as vaginal exposure that is initially treated with local excision and primary closure might present with dyspareunia or recurrent exposure long after the follow-up period has expired.^{9,31,41–44} Indeed, many patients in one of our own studies presented in exactly this way, but this fact was never captured in the paper owing to the methodology used.⁹ We have seen many unreported examples of patients managed conservatively with short-term success, who ultimately presented with a recurrent complication that occurred after the study ended, owing to the short follow-up duration of most published research (1–2 years) relative to the expected lifespan of most implanted slings.^{9,81,105,143–147} Authors of one study estimated the overall incidence of vaginal extrusion of mesh and pelvic pain to be 6% and 4.3%, respectively.⁴⁵ Despite peer-reviewed literature in this area being replete with statements about the short-lived nature of sling-related complications, most case reports of mesh sling complications document the treatment-refractory nature and suboptimal outcomes associated with these complications, none of which was captured by the original studies.^{9,43} For some complications (cystitis, voiding dysfunction, pain or neurological symptoms), most authors claim that only expectant or medical treatment is necessary and that patient outcomes are satisfactory, without presenting any meaningful follow-up data to substantiate these claims.^{30,146,148,149}

In 2011, the International Urogynaecology Association (IUGA) and the International Continence Society (ICS) published a joint recommendation for a standardization of terminology to report complications related to the insertion of prostheses and grafts in female pelvic floor surgery.¹⁵⁰ To date, these guidelines have not been widely used. The net result of all of this is that the science of assessing and reporting midurethral sling complications is seriously flawed.

Table 5 | Comparison of complications of patients with an RP sling or TOT sling

Complication	RP sling (mean; range)	TOT sling (mean; range)	Combined RP and TOT sling (mean; range)
General complications			
Death within 30 days	0.0	0.0	0.0
Urethral obstruction/voiding dysfunction	8.8; 0–32.7	5.9; 0–33.9	7.3; 0–33.9
Requiring surgery	2.7; 0–8.9	2.0; 0–21.3	2.3; 0–21.3
Urinary infections	8.6; 0–39.1	6.2; 0–23.3	7.3; 0–39.1
Pain (within 6 weeks)	4.5; 0–23.1	10.2; 0–33.3	7.9; 0–33.3
Neurologic symptoms (within 6 weeks)	1.6; 0–5.0	0.9; 0–10.3	1.2; 0–10.3
<i>De novo</i> OAB	11.4; 0–29.4	10.3; 0–48.1	10.9; 0–48.1
Pelvic organ perforation			
In total	4.8; 0–14.3	2.3; 0–16.1	3.5; 0–16.1
Bladder	4.6; 0–14.3	1.1; 0–16.1	2.9; 0–16.1
Vaginal	1.0; 0–2.1	1.6; 0–14.1	1.4; 0–14.1
Urethral	0.0; 0–1.5	0.0; 0–0.3	0.0; 0–1.5
Bowel	0.0; 0–1.7	0.0	0.0; 0–1.7
Ureteral	0.0	0.0; 0–1.3	0.0; 0–1.3
Mesh exposure/erosion/extrusion			
In total	2.3; 0–26.1	2.7; 0–19.0	2.5; 0–26.1
Treated conservatively	0.1; 0–5.6	1.0; 0–7.1	0.9; 0–7.1
Vaginal	0.0; 0–4.6	1.0; 0–7.1	0.7; 0–7.1
Bladder	0.0; 0–5.6	0.0	0.0; 0–5.6
Urethral	0.0	0.0	0.0
Requiring surgery	1.6; 0–26.1	1.9; 0–15.9	1.8; 0–26.1
Vaginal	1.0; 0–10.9	1.8; 0–15.9	1.5; 0–15.9
Bladder	1.4; 0–15.2	0.0*	0.2; 0–15.2
Urethral	0.3; 0–16.7	0.0; 0–2.6	0.2; 0–16.7
Longer-term complications			
Refractory pain (>6 weeks)	2.0; 0–7.9	5.3; 0–30.5	4.1; 0–30.5
Neurological symptoms (>6 weeks)	1.0; 0–5.2	0.9; 0–10.6	1.0; 0–10.6
Fistulas	0.4; 0–0.7	0.2; 0–1.1	0.3; 0–1.1

*Case series only investigating bladder erosions were removed from the analysis. Abbreviations: OAB, overactive bladder; RP, retropubic tension-free transvaginal mesh tape; TOT, transobturator tape.

Infection

In a randomized study of SMUS effectiveness, investigators found culture-proven cystitis in 8.4% of patients with RP and 4.7% with TOT slings.³⁰ In another study, the authors reported that 12% of patients developed at least one UTI in the first 3 months after RP sling surgery.¹⁵¹ Our literature search documented bacterial cystitis in 0–39% of patients who underwent SMUS surgery (REFS 13,18,21–23,30,39,47,48,50,53,55–57, 59,61,66,74,76,83,91,152–186). Unfortunately, limited published data are available on the long-term consequences of these infections. For example, in one study, 7.3% of women had recurrent UTIs 12 months after being fitted with a TOT sling, but no published reports of longer-term follow-up currently exist.⁷⁶

Other more serious infections have been reported after SMUS implantation. In a comprehensive literature review⁴⁵ several such complications following sling placement were reported: cellulitis,^{117–121,187} abscess formation, including in the retropubic space,¹²² retroperitoneal space,¹²³ thigh,¹²³ obturator space^{124–126} and ischiorectal fossa;^{127,128} sinus tract formation;¹²⁹ necrotizing fasciitis;¹³⁰ osteitis pubis;¹³¹ and sepsis.¹² Thigh abscesses, a complication unique to the TOT sling approach, have been reported in the past decade.^{101,123,188,189} Occurrence of such serious SMUS-related infectious complications is often delayed by months or years after sling implantation. Presenting symptoms include chronic discharge from the vagina, thigh and/or perineum.

Treatment of infected mesh and abscesses requires open drainage and removal of all mesh, which, otherwise, will serve as a nidus for further infection. Amid types II and III mesh materials,¹⁹⁰ such as Silastic® (Dow Corning, MI, USA) and Gore-Tex® (W.L. Gore & Associates, DE, USA), are easily identified and pulled out, usually in their intact form. The technical challenges of removing type I meshes are considerably greater. For example, even though the entire mesh is likely to be infected, only part of it is involved in the abscess. This part is easy to remove, but because of tissue ingrowth and probable degradation, the remaining mesh is usually embedded in the tissue and might fragment during dissection. Furthermore, the retropubic and thigh portions of RP and TOT slings, respectively, are notoriously difficult to remove.^{9,43,78,84,85,94,100,138} RP slings are often adherent to the bladder neck and difficult to access during surgery. We, and others, have noted that the infection can track along muscle planes and even form a psoas muscle abscess in some patients.^{123,191} These abscesses might require multiple operations in order to achieve a satisfactory outcome. Unfortunately, owing to the technical reasons explained above and concerns about complications in adjacent organs during the dissection, complete mesh removal is often not accomplished in patients with type I mesh slings. Necrotizing fasciitis (Fournier gangrene) has also been reported after both RP¹⁹² and TOT¹⁹³ sling implantations. In patients with SMUS-related infections, removal of the complete mesh is particularly important. Of course, many more serious infectious complications are likely to arise from the ‘minor’ ones listed above,^{113,126} but we could find no published studies that actually address this issue.

Pelvic organ perforations

Pelvic organ (bladder, urethral, vaginal [REFS 10,14, 16,18,21–23,30,33,39,47,48,50,52,53,55,56,59,62–64, 66,67,69–71,73,74,76,77,80,81,86–88,90–93,98, 99,109,132–134,140,145,146,148,151–153,156,158, 162,163,167,168,171,173,176–179,181–184,194–238], or bowel) perforation at the time of trocar passage has been reported to occur in 0–16% of sling surgery procedures (0–14% for RP sling implantation and 0–16% for TOT sling implantation). In most reports, the authors usually downplay any substantial implications of pelvic organ perforations. Authors will typically state that

Box 1 | Summary of mesh safety**Complications requiring surgery**

- Urethral obstruction 3.2%
- Erosion, extrusion or exposure 2.0%
- Fistulas 0.3%
- Bowel injury or infection 0.1%
- Lifestyle-altering complications
- Chronic pain 4.3% (0.5%)*
- Refractory (*de novo*) OAB 11% (3.9%)*
- Recurrent and/or persistent SUI 5.3%
- Total incidence of serious complications and/or sling failure in patients with incontinence 15.3%

*Indicates numbers are not mutually exclusive to individual complications. For example, a patient who underwent mesh excision for exposure may develop refractory pain. The numbers in parenthesis refer to the estimated incidence of complications refractory to medical or surgical treatment. Abbreviations: OAB, overactive bladder; SUI, stress urinary incontinence.

recognized trocar perforations of the bladder might be treated by simply removing the trocar (or sling) and passing it again and, at the discretion of the surgeon, leaving an indwelling catheter in for a matter of days or, sometimes, not at all.^{148,239–242} However, in an article published in 2014, the authors concluded that occurrence of bladder and/or urethral perforations during surgery is associated with an almost 26-fold increase in risk of subsequent bladder or urethral mesh erosion.³¹ If the findings of this study are accurate, the dictum of simply removing and repassing the trocar after bladder perforation must be seriously questioned.

Limited information is available on urethral complications following perforation, but if a perforation large enough to necessitate repair exists, most would agree that it is best to abandon the SMUS procedure.^{39,62,64,73,140,213} Depending on the circumstances, buttressing the repair with a Martius flap in anticipation of performing an autologous or allograft sling procedure at another date is one possible approach; alternatively, in rare circumstances, when it is desirable to complete the surgery in one sitting, this could be done at the same time as sling implantation.^{9,43,243} Such a situation might arise when a patient's health dictates that the risk:benefit ratio favours a single operation instead of two or, if the patient lives at a great distance from the treatment centre and returning for a second operation would present a major burden.

Bowel perforations during sling surgery are even more uncommon than perforations of other pelvic organs with a reported incidence in 0.005–0.02% of procedures.¹³² In a meta-analysis published in 2007, authors reported a mortality rate of 20% in 35 incidences of bowel perforation during sling surgery.¹³³ Of particular concern is the fact that some patients did not present with any clinical signs of abdominal injury.^{102,104,106,132–134,236} Patients' initial complaints were often originally attributed to voiding dysfunction and not until days later, when the patients became seriously ill, was the correct diagnosis made.²³⁷ In five patients the diagnosis of bowel perforation during sling surgery was made only at autopsy.²³⁸

Voiding dysfunction

The term 'voiding dysfunction' was not clearly defined in the majority of studies in this area and could be interpreted as either voiding symptoms, storage symptoms or both. Some of the terms used include *de novo* or persistent overactive bladder, urgency or urge incontinence, urinary retention and urethral obstruction from SMUS. In addition, secondary voiding problems can arise after sling revision or excision surgeries performed to treat the original complications such as recurrent SUI, dyspareunia, urethral strictures and fistulas.^{41,100,111,244} We used our best judgement to assign the authors' intent to one of the categories listed below, but in some instances the author's intent was not clear. For example, one category was listed as "voiding dysfunction requiring surgery",³⁰ although whether this was a result of urethral obstruction or refractory OAB was not defined. Keeping in mind the caveats listed above, temporary or refractory voiding dysfunction has been reported in 0–33% of patients with a SMUS, and is more common in patients fitted with an RP sling than in those fitted with a TOT sling (REFS 16–23,26,30,33,39,46,47,49–51, 53,54,56,58,59,61–66,68–71,73,74,76,77,79,80,82, 83,86–93,95,97–99,140,145,146,148,151–153,156–164, 166–171,173,174,176–184,186,194–205,207,208, 211,214–218,220,222–229,231–233,235,245–260).

OAB symptoms

The terms used to describe OAB in the studies reviewed included urge or urgency incontinence, urgency, refractory urgency and overactive bladder.^{9,261} In addition, the qualifiers persistent or *de novo*²⁶¹ were often used. *De novo* OAB, indicating the occurrence of OAB after sling surgery, was reported in 0–48% of patients in various studies (REFS 17,18,20,26,39,46–48,50–53, 55,57–59,61,62,66,67,70,73–77,79,83,87,88,91,92, 96,97,99,140,152,153,155,157,159,161,163,164,166–168, 170–174,177,178,181–186,209,211,213,216–219,222, 224,228,229,231–235,245,249,254,255,259,262–269). Most patients with OAB had symptoms that were said to have resolved within the first month of surgery either spontaneously, or in response to anticholinergics, antibiotics or self-catheterization.^{30,194} However, the metrics used to conclude this were inadequate for the task; in fact, the vast majority of studies used no metrics or validated outcome measures at all. When validated instruments were used, they were often, in our judgement, inadequate. For example, the UDI was one of the most common questionnaires used and, as alluded to above, conflates the degree of incontinence with bother.²⁴ Furthermore, the UDI contains no question that specifically refers to urgency (as opposed to urge incontinence). Approximately two-thirds of women with OAB do not have urge incontinence;²⁷⁰ thus, use of this instrument is likely to miss two-thirds of the women with urgency or OAB in any series.

In patients with refractory OAB as a SMUS-related complication, a careful search for a remediable underlying aetiology should be conducted. Possible aetiologies include infection, stones, urethral obstruction

and mesh erosion into the bladder or urethra. When such aetiologies are found and treated, the reported success rates are generally high, but the methods used to determine treatment success were generally of poor quality. For example, in reports from two studies of endoscopic laser ablation of eroded mesh in patients with OAB the authors reported successful outcomes, but did not use any objective outcome measures.^{100,108} Other studies did use validated instruments to quantify outcomes such as the Overactive Bladder Symptom Score, voiding diaries and pad-weight tests.⁹ With these caveats in mind, a successful outcome after surgical treatment of these remediable conditions was reported in 28–64% of patients with refractory OAB as a SMUS-related complication.^{9,30,43,85}

Urethral obstruction

Urethral obstruction is a urodynamic diagnosis based on high detrusor pressure accompanied by low urine flow rate, although again, no uniform criteria for diagnosing obstruction were used in the papers reviewed. Most investigators simply inferred obstruction based on the temporal relationship between SMUS surgery and voiding symptoms.^{271,272} Others used measurements of urine flow rate and post-void residual volume or urodynamics. Urethral obstruction should be suspected in any woman with persistent voiding symptoms (either storage or emptying) after SMUS placement.^{9,94,273,274} Obstruction is definitively diagnosed by the findings of high detrusor pressure and low uroflow during urodynamics. Generally, the existence of a normal urine flow rate is thought to exclude the presence of urethral obstruction; however, this is not always the case, as sometimes abnormal urine flow rate can be generated by a strong detrusor contraction or abdominal straining (Figure 1).

Even in the absence of urodynamically confirmed urethral obstruction, sling incision and/or excision can completely resolve refractory voiding (and OAB) symptoms. Compression from the sling is by far the most common cause of urethral obstruction, and at least one case of urethral stricture accompanied by urethral erosion has been reported.⁹ The incidence of urethral obstruction requiring surgical intervention ranges from 0% to 8.9% in patients fitted with RP slings and from 0% to 21.3% for those with TOT slings (REFS 17,18,20,21,23,26,30,46–48, 50–56,58,59,61–64,66–69,71,74,77,80–83,86–92,97–99, 148,159,163,166,173,178,180–182,216–218,233,245,250, 253,255,258–260,269). In the largest patient series reported to date, which comprised nearly 190,000 SMUS implantation procedures, and was based on insurance data, a 9-year cumulative rate of sling revision surgery owing to urethral obstruction of 1.3% was reported. Most of this revision surgery occurred in the first few years after the original implantation procedure.⁶ In this patient series, however, no mention of the number of patients lost to follow-up monitoring was made, and because investigators only searched for one Current Procedural Terminology (CPT) code (57287) for the sling revision surgery, the actual incidence of revision surgery owing

to urethral obstruction was possibly higher. Similarly, a rate of urethrolysis of 3.4% was reported in a cohort of 818 patients fitted with RP slings, although investigators failed to search for ‘mesh removal’, ‘explant’ or ‘excision’.²⁷⁵ Owing to incomplete use of these search terms, retrospective studies of databases^{6,270} are prone to underestimating the incidence of urethral obstruction requiring surgery.

The reported incidences of urethral obstruction requiring surgery are generally well under 10%. A small number of patients having urethral obstruction in the lost-to-follow-up group could substantially increase the actual overall incidence. In fact, most studies found that 50–75% (sometimes more) of patients undergoing sling revision surgery were treated by a surgeon other than the implanting surgeon.^{9,44,276}

Some authors have recommended sling incision or even urethral dilatation for treatment of patients with urethral obstruction, although most authors agree that the entire suburethral portion of the sling should be removed, even if an incision into the wall or urethral lumen is required.^{9,40,154,244,261} Whether to remove all of the mesh from RP slings in patients with urethral obstruction depends on multiple factors, including associated pelvic pain, dyspareunia and/or recurrent infections that might be related to retained mesh.^{277–280} No meaningful data exist regarding the effectiveness of urethral dilatation; however, based on our clinical experience, we believe that this approach should not be used owing to the possibility of a urethral abrasion that might ultimately lead to erosion.²⁸¹ Of course, optical urethrotomy, internal urethrotomy and transurethral incision of urethral strictures should not be done at all except in the rarest of circumstances, for fear of causing iatrogenic urethral exposure.

No clear indications for urethrolysis currently exist; rather, the need for procedures of this type should be considered on a patient-by-patient basis, depending on the degree of scarring and urethral immobility.^{282,283} In our judgement, urethrolysis should be considered in patients in whom the proximal urethra feels scarred and immobile during surgery and/or following a finding of limited urethral mobility on a Q-Tip® (Unilever, London, UK) test.

A high, and well documented incidence of recurrent SUI after mesh revision surgery exists, that is reported to range from 10% to 60% of patients who undergo revision surgery.^{9,43,78,84,85,94,100,138} For each patient with symptoms of recurrent sphincteric incontinence, a decision needs to be made as to whether or not another anti-incontinence procedure should be considered. Reports of sling revision surgery in patients with sphincteric incontinence are sparse, and often anecdotal; but, if urethral reconstruction is necessary at the time of mesh removal, the AUA guidelines on the surgical management of female SUI¹ state that implantation of another SMUS are contraindicated in these patients. Most authors recommend a wait-and-see approach to management of patients with recurrent SUI after mesh revision surgery, and, as a rule, we agree with this




Figure 1 | Identification of urethral obstruction. **a** | Urodynamic trace showing severe urethral obstruction (Blaivas–Groutz nomogram type 2) caused by the presence of a urethral stricture in a 52 year-old woman 4 years postimplantation of a SPARC™ (American Medical Systems, MN, USA) SMUS. Urethral obstruction is confirmed by the presence of strong, sustained detrusor contraction ($P_{det} Q_{max} = 56$ cm H_2O and $Q_{max} = 1$ mL/s). **b** | Cystourethrogram confirming the presence of an obstruction owing to the presence of a midurethral stricture (arrows). **c** | Urodynamic trace showing high-flow urethral obstruction in a 52 year-old woman 6 years after midurethral SMUS implantation. Urethral obstruction was confirmed by $P_{det} Q_{max} = 56$ cm H_2O (Blaivas–Groutz nomogram type 1). Owing to a technical error, infused volume was not recorded on this trace. **d** | Voiding cystourethrogram showing the site of urethral obstruction to be in the distal third of the urethra (arrows). Abbreviations: EMG, electromyogram; P_{abd} , abdominal pressure; P_{det} , detrusor pressure; $P_{det} Q_{max}$, voiding pressure at peak flow; P_{ves} , intravesical pressure; Q_{max} , peak flow; SMUS, synthetic midurethral sling; V_{H_2O} , bladder filling volume.

recommendation. However, data from two published reports contradict this recommendation. A continence rate of 71% was reported in 14 patients who underwent mesh removal after urethral perforations and, in another study, an 82% success rate after revision surgery was reported in 28 patients, of whom 14 had a synchronous autologous sling. In both series a Martius flap was placed between the urethra and sling.^{9,43,243} No clear indications exist for conducting synchronous anti-incontinence surgical procedures during mesh excision surgery, although we believe that this approach should be considered whenever extensive damage to the proximal half of the urethra has occurred. Authors of most study reports, however, do not report thoroughly on the outcomes of sling excision surgery.

Mesh erosion, extrusion or exposure

Reports of research in this area of sling complications are typically replete with terminology that conflates the terms erosion, extrusion and exposure; thus, discerning the exact meaning of every author was often impossible. The joint recommendations of the IUGA and ICS¹⁵⁰ provide guidance regarding use of terminology related to SMUS complications, but, in a clinical sense, applying the recommended distinctions is usually not possible in most patients. The IUGA and ICS guidelines define exposure as “a condition of displaying (mesh), revealing, exhibiting, or making accessible for example through the vagina”¹⁵⁰ and extrusion of mesh as “passage gradually out of a body structure or tissue.”¹⁵⁰ The guidelines also recommend avoiding use of the term ‘erosion’ altogether,

but in this Review, we use the terms interchangeably as there is no sound scientific way of making this distinction exists given that the overwhelming majority of authors use this terminology interchangeably. Conceptually, mesh can be seen to protrude through the vaginal wall or into the bladder, urethra or bowel by one of two mechanisms: either it was inadvertently positioned there at the time of surgery or somehow, over the course of time, the mesh gradually worked itself into such a position.

The incidence of mesh sling erosions varies widely between study reports, ranging from 0% to 41% of patients (REFS 16–20,22,26,30,31,39,46–50,52,53, 55–73,75–77,79,81,82,86–90,92,93,95–100,152,153, 155–160,162,164,166–179,181–185,204,207–209, 212–215,217–221,224,225,229,231,233–235,245,247, 252–255,257–259,265,266,268,284–293). The risk factors for sling erosion fall into three main categories: patient factors; mesh characteristics; and intraoperative considerations.⁴² With respect to the patient, oestrogen-deficient states, genital atrophy, surgical scarring, concurrent prolapse surgery, type 1 or type 2 diabetes mellitus, steroid use, concurrent anticholinergic use and smoking have been reported as risk factors for sling erosion.²⁴⁶ Patients ≥ 75 years of age also had a higher incidence of OAB and recurrent UTI²⁶⁶ and patients of both younger and older ages (mean ages 55, and 75 years respectively) were variously reported as having adjusted risk factors. Previous pelvic radiation is another obvious risk factor for mesh erosion, but few of these patients undergo sling surgery; thus, this factor did not appear as such in the literature.

Certain types of mesh have a particularly high risk of erosion based on the intrinsic characteristics of the materials they are made from. In a study published in 1997,¹⁹⁰ synthetic materials used for herniorrhaphy (a type of hernia repair surgery) were categorized based on their composition (synthetic or biological), structure (monofilament or multifilament), pore size (macroporous or microporous) and architecture (knitted or woven). Type I (knitted, monofilament and macroporous polypropylene mesh) is currently considered to be the optimal SMUS mesh material owing to its large pore size ($>75\ \mu\text{m}$), which facilitates infiltration of macrophages and fibroblasts, promotes neovascularity and tissue ingrowth, and minimizes the likelihood of infection. Examples of Type I mesh include VitaMESH™ (Atrium, NH, USA), Marlex® (C.R. Bard, NJ, USA), Prolene® (Ethicon, NJ, USA) and Trelex Natural® mesh (Boston Scientific, MA, USA).

In an attempt to decrease the foreign body responses associated with mesh materials and increase tissue compliance, several manufacturers have designed lightweight meshes of decreased density with smaller fibre diameter and larger pores, with the intention of preventing stiffness, contraction and mesh shrinkage. Several published studies purport some benefit of these new materials in patients requiring inguinal hernia repair; however, all of the studies involved small numbers of patients, with limited follow-up duration, thus precluding any meaningful conclusions.^{35,294–300}


Amid type II mesh (monofilament and microporous) has pores ($<10\ \mu\text{m}$ in diameter) that are large enough to allow bacterial infiltration but too small for macrophage infiltration, thus infection is more probable and tissue ingrowth is impeded.¹⁹⁰ Polytetrafluoroethylene (PTFE, Gore-Tex® W.L. Gore & Associates, DE, USA) is the most common prototype type II mesh. Surgical Membrane Type III multifilament mesh is much denser and stiffer than other types of mesh and has interstices that are $<10\ \mu\text{m}$ in diameter with the same negative consequences as those of type II mesh. PTFE mesh (Teflon® DuPont, DE, USA), braided polyethylene terephthalate mesh (Mersilene® Ethicon, NJ, USA), braided polypropylene mesh (Surgipro™ [Covidien, CA, USA] monofilament mesh) and perforated PTFE patch (GORE® MYCROMESH® W.L. Gore & Associates, DE, USA) are examples of type III meshes. Type IV meshes are submicroporous coated biomaterials with pores $<1\ \mu\text{m}$ in diameter. SILASTIC® (Dow Corning, MI, USA), Celgard® polypropylene sheeting (Celgard, NC, USA) GORE® PRECLUDE® Pericardial membrane and GORE® PRECLUDE® Dura-substitute (both manufactured by W.L. Gore & Associates, DE, USA) are all type IV meshes. Types II–IV meshes, including PTFE mesh (Amid Type II), silicon-coated polyethylene or polyester (Amid Type IV) and non-knitted, nonwoven mesh such as OBTAPE® and UraTape® (both Mentor–Porges, Le Plessis Robinson, France), have been documented to have a much higher incidence of erosion (16–25%) compared with that of type I meshes (0–10%).^{105,109,124,301}

As described previously, bladder, urethral or vaginal perforation during the original surgery increases the risk of subsequent sling erosion by approximately 26-fold.³¹ In addition, passage of the trocar through the vaginal, bladder or urethral wall without actually penetrating the lumen might occur, which would result in positioning of the mesh just barely under the surface of the lumen and predisposing it to erosion. This possibility seems likely, although it is currently unproven.

The approach to treatment of sling erosions has been largely empirical, follow-up durations of studies in this area have been short and only a few studies have applied validated outcome measures, especially with respect to the occurrence of other symptoms, such as lower urinary tract symptoms, pain and dyspareunia.^{9,44} As discussed previously, recurrent SUI after mesh removal is not uncommon.^{9,43,78,84,85,94,100,138} However, synchronous anti-incontinence surgery can be effective in this setting.⁴³ The authors of this study⁴⁰ based their decision to conduct synchronous autologous sling surgery on multiple factors: the location of the urethral injury; preoperative continence status; and the degree of urethral hypermobility. In our series,⁹ the success rate (based on PGII score) was 82% after mesh removal, but only half of the patients underwent synchronous autologous sling surgery.⁹

Mesh erosion in the bladder

Bladder erosions are reported to occur in 0–15% of patients fitted with a SMUS (REFS 16–20,22,26,30,46–48,



◀ **Figure 2** | Identification and removal of eroded mesh.

a | Eroded mesh can be difficult to see during cystoscopy owing to its almost translucent appearance in some patients. **b** | In this erosion, the sling is nearly obscured by the presence of calcium deposits. **c** | Appearance of eroded Amid Type I mesh at urethroscopy. This mesh had no appreciable tissue ingrowth and pulled out of the urethra easily, leaving a small urethrotomy that was closed with a few sutures. **d** | Surgical explantation of eroded silicone mesh. In this case, removing the mesh was relatively easy because it was an Amid type IV mesh, which was encapsulated. **e** | Transvesical explantation of an eroded mesh sling. This type I mesh had tissue ingrowth and required delicate surgery and sharp dissection to enable removal. **f** | Following sharp dissection, the mesh was completely excised from the bladder wall. It coursed over, but did not damage, the ureter. Permission obtained from Fred Govier and Kathleen Kobashi, Virginia Mason Hospital & Seattle Medical Center, Seattle, WA, USA.

is necessary to remove all the intravesical mesh;^{9,10,313} laparoscopic approaches have also been tried.¹³⁸ Unfortunately, few studies of patients with mesh erosions in the bladder have sufficiently long follow-up durations or good enough outcome measures to determine the true success rates of these surgeries. Furthermore, all reported successes and failures were compared to the patient's status before the mesh removal surgery, and not before SMUS implantation surgery.

Mesh erosions in the urethra

Urethral erosions are much less common than bladder erosions in patients fitted with a SMUS, with a reported incidence of 0–2.6%.^{46,61,100,176} Urethral strictures caused by mesh erosions are even rarer than urethral erosions.³¹⁴ A number of different aetiologies of urethral erosion have been postulated, including surgical transgressions (excessive sling tension, unrecognized urethral perforation and passage of the sling through the urethral wall), urethral dilatation,²⁸¹ postoperative contraction of the sling,³¹⁵ infection, inflammation and immunological reactions.^{33,315–319} Treatment of urethral erosions might be as simple as excision of the mesh, which sometimes pulls easily out of the urethra, (Figure 2) or can involve extensive surgical excision of the mesh with part of the urethral wall requiring urethral reconstruction and a Martius flap (Figure 3).³²⁰

Vaginal mesh erosion, extrusion or exposure

Vaginal mesh erosion, extrusion or exposure has been reported in 0–19% of patients with a SMUS (0–11% of patients with an RP sling and 0–19% of patients with a TOT sling, REFS 16–20,22,26,30,31,46,48,50,53,58–60, 62,64,65,67,69,70,72,75–77,79,81,82,86–90,92, 95,97,103,140,156,158,160,164,167,170,175,177,183, 185,218,221,231,233,247,257,260,266,288,291,292). Factors associated with a higher incidence of mesh extrusion include trocar perforation of the vaginal wall during mesh implantation, previous pelvic surgery, diabetes, bleeding complications at the time of surgery, pelvic radiation, smoking, older age and vaginal incision length >2 cm.^{31,42,321}

50,52,53,55–73,75–77, 79,81,82,86–90,92,93,95–100, 140,152,155–160,164,166–168,170–172,174–177,181, 183–185,207–209,213,215,217–219,221,229,231,233, 234,245,247,253,254,257,258,260,265,266,288–293). Patients with mesh erosions in the bladder usually present with recurrent UTIs, haematuria, bladder stones, incontinence, dyspareunia and pelvic pain; mesh erosions are typically discovered during cystoscopic examinations (Figure 2). Most authors agree on the general treatment principle—all mesh must be removed from the bladder—but the procedures used to do so vary widely. A variety of endoscopic approaches have been used to remove mesh including cutting with scissors and removing the mesh with grasping forceps, transurethral resection of the mesh using monopolar or bipolar current, vaporizing it with a holmium laser^{302–310} and even utilizing a small nasal speculum or Metzenbaum scissors passed transurethrally alongside a cystoscope.³¹¹ Results achieved with these surgeries^{300–309} have been variable, with some successes followed up for as long as a few years, but the majority of studies had short follow-up durations. Open surgery using a suprapubic or vaginal approach has the advantage of removing all of the intravesical mesh, including mesh that traverses the bladder wall (Figure 2).^{9,43,312} Sometimes, partial cystectomy




Figure 3 | Identification and removal of Amid type I eroded mesh surrounding the urethra. **a** | Urethrosopic view of urethral erosion. In this case the erosion was very subtle and located in the distal urethra, such erosions are very easy to miss. **b** | Transvaginal dissection and isolation of the mesh shown in image a reveals marked scarring and tissue ingrowth. **c** | Removal required sharp dissection that left a large defect in the urethra (arrow) that required urethral reconstruction, Martius flap and implantation of an autologous fascial sling. **d** | The sling being placed over the urethra. **e** | Creation of a Martius flap (arrow) between the reconstructed urethra and sling, thus repair is completed before vaginal wound closure.

We postulate, based on our clinical experience, several other causes of vaginal mesh exposure including wound dehiscences resulting in exposure of the mesh or implantation of the mesh superficial to the pubocervical fascia so that it lies just beneath the surface.⁹ When conditions that favour mesh erosions in the vagina are compounded by local ischaemia, inflammation, foreign body reaction and/or infection, the risk of erosion is likely to be increased. An alternative causative factor has also been suggested—defective wound healing caused by an immunological response to the mesh itself.³³ Vaginal mesh extrusion frequently presents as dyspareunia, vaginal discharge, vaginal bleeding and pain experienced by the sexual partner during vaginal intercourse (“hispareunia”).³²² Sometimes asymptomatic extrusions are found during a routine vaginal examination. A diagnosis of vaginal mesh extrusion is typically based on a physical examination—by visual inspection and/or palpation (Figure 4). Most vaginal mesh extrusions occur within the first year of SMUS implantation, although they have been observed in patients as long as 17 years after the original surgery.¹⁶ Some authors report that small areas of mesh exposure can be successfully treated with topical oestrogen,³²³ although the results have been mixed.³²⁴ Larger mesh

exposures require primary closure of the vaginal wall over the exposed mesh or surgical excision and closure with or without vaginal wall flaps.^{30,41–43,45} Mesh exposures of this type are usually reported as minor complications; post-treatment follow up in these patients has been woefully inadequate in most studies.^{319,323–325} We did not find sufficient justification in the literature to be able to confidently assess the long-term success of treating these minor exposures. For example, in a retrospective review of nearly 347 complications, the authors found that 73% of patients who initially had nonsurgical treatment for vaginal mesh extrusions ultimately required surgical treatment within 5 years of the original sling surgery.^{16,44} In a single-institution study of 79 patients who underwent SMUS implantation, the mean time from SMUS implantation to removal was 2 years with a range of 0–11 years. Despite the fact that mesh erosions can occur >10 years after implantation, most studies report a mean follow-up duration of only 2–23 months.^{16,105,261,277,301,312,326}

Mesh erosions in the bowel

Bowel mesh erosions are exceedingly rare and all known patients with bowel erosions have presented with an enterovaginal fistula.^{9,327,328} Treatment requires removal of all intra-abdominal mesh and repair or excision of the affected bowel. In our own clinical experience, a patient who had bowel mesh erosion had an associated vesico-vaginal fistula and after several unsuccessful reconstructive surgeries required a continent urinary diversion.⁹

Pain

Pain is the most poorly studied complication of SMUS surgery; we found only a few studies that included prospective data collection and/or validated questionnaires assessing pain.^{30,79,90,95,152,168,174,229,254} Some investigators used postoperative questionnaires that contained pain questions, but most relied on patients' recall or chart review.³⁹ In addition, many different descriptors of pain have been reported including vaginal, pelvic, groin, thigh, leg, suprapubic and lower abdominal pain, dyspareunia and “pain, patient self-report.”³⁰ Two different kinds of pain caused by nerve injury have been suggested: centrally mediated hyperalgesia and a peripherally mediated painful hypoalgesia, suggesting the need for mechanism-based classification of neuropathic pain.³²⁹ Most importantly, few of the case studies of patients with SMUS-related pain quantified the severity of this pain,




Figure 4 | Vaginal complications and removal of SMUS. **a** | View of mesh extrusion into the vagina. This extrusion was both readily visible and appeared palpable on physical examination. **b** | Transvaginal view of eroded mesh obscured by a granuloma. The erosion was neither visible nor palpable. **c** | Transvaginal view of fistula obscured by granuloma. **d** | Transvaginal explantation of an Amid type I RP sling. Note the dense tissue ingrowth and scar that requires sharp dissection in order to remove the suburethral portion in its entirety. This portion of the sling felt thickened and stiff. **e** | Retropubic view in the same patient, as shown in image d. The retropubic portion of the mesh after dissection showed fatty tissue ingrowth. This portion was pliable and not scarred. Abbreviations: RP, retropubic; SMUS, synthetic mid-urethral sling.

its character or how the pain affected patients' quality of life.^{30,64,82,263} This neglected topic is of the utmost importance and permanently affects "a small cohort of patients whose lives have been unalterably changed for the worse."^{40,312}

Pain in patients with a SMUS has been attributed to direct nerve injury during implantation, nerve entrapment, haematoma, infection, chronic inflammation, structural changes to the implanted mesh (shrinkage, stiffening, hardening and/or banding) and scarring.^{38,330} Most patients present with pain within the first year of surgery, although some present years later—as late as 8 years postoperatively.^{9,110,143} In studies of effectiveness and safety, pain was mostly divided into perioperative pain and pain lasting more than 6 weeks. Perioperative pain has been reported in up to 33% of patients,²⁵² occurring more frequently after implantation of TOT slings than RP slings,³³¹ and chronic pain (of any definition) has been reported in 0–31% of patients (REFS 17,18,20,30,39,47,48,53,58,59, 61–63,66,69,70,73,76,79,82,89–91,95,98,99,140,152,155, 159,164,167,168,171–174,177,180–182,185,208,209, 215,223,229,231–233,235,247,250,252,254,255,257,266, 268,290,332,333).

Chronic disabling pain is one of the most common indications for mesh removal, particularly in patients fitted with TOT slings.^{9,38,42,44,110,111,277,326} Chronic pelvic pain often contributes to a need for mesh removal; however, these data might not be captured by the approaches used by all investigators. Thus, the reported incidence of pain complications is likely to be falsely low. In comparison with patients with an RP sling, patients with a TOT sling have a higher incidence of persistent pain (32% versus 10%) and dyspareunia (18% versus 3%).¹¹⁶ This finding is confirmed by a review and meta-analysis in which the rates of chronic groin and leg pain were higher in patients with a TOT sling compared with those of patients with an RP sling (16% versus 6.5%, respectively).²⁰⁶

Treatment of persistent pain in patients with a SMUS is particularly challenging and has been entirely empirical and progressive in nature. The treatment approach in these patients typically begins with pain medications and neuromodulatory medications such as carbamazepine, physical therapy or trigger-point injections and culminates with partial or complete mesh excision. Reported success rates of these treatments range from 24% to 100%,^{9,40,43,107,277,312,326} but use of validated outcome measures documenting treatment success and long-term follow-up monitoring are both lacking. Furthermore, a number of case studies and series of patients with mesh complications have commented on the lifestyle-altering nature of painful complications in these patients.^{9,40,107,312,326}

Fistulas

Urethrovaginal and vesicovaginal fistulas are rare SMUS-related complications, with a reported incidence of <1%.^{46,48,66,98,173,178,212,217,334} These fistulas are most frequently associated with bladder or urethral erosion of the sling and patients can present in a variety of ways, and as late as 6 years after the initial surgery.^{41,43,44,335–338} Despite their low reported incidence, the possibility of fistulas should be considered when patients present with recurrent incontinence, OAB, pain and/or voiding dysfunction after mesh surgery.⁴¹ Not infrequently, a diagnosis of fistula can be obscured, owing to the presence of adjacent granulation tissue (Figure 4). Concurrent sphincteric incontinence might also confound diagnoses of fistulas,

especially in patients with a urethrovaginal fistula; thus, a careful evaluation should be undertaken to exclude fistula whenever a patient has recurrent incontinence after mesh sling surgery. This evaluation should include a physical examination with a stress test and visual confirmation that leakage is occurring through the urethral meatus as well as cystourethroscopy.⁴¹ Surgical repair of urethral or vesicovaginal fistula requires the complete removal of all involved mesh and possible vaginal reconstruction with tissue flaps.³³⁹ In patients with concomitant sphincteric incontinence, synchronous repair of a urethrovaginal fistula and an autologous fascial sling with a Martius flap interposed between the fistula repair and sling is often an effective treatment.^{41,43}

Death

Mortality is the least common SMUS-related complication. In fact, in our literature review we did not find a single case series report that contains a postoperative death of a patient undergoing SMUS implantation surgery; However, In a study of bowel complications of SMUS, published in 2007,¹³³ 7 deaths from bowel injuries after RP sling implantation were reported, and in 2014 another death after bowel perforation during a retropubic SMUS implantation was published as a case report.¹³² Authors of a review of database entries regarding SMUS complications reported 10 deaths owing to bowel injury (six), vascular injury (three) and sepsis (one).¹⁰ Authors of this study¹⁰ suggested that death is an under-reported complication in patients treated with a SMUS.

Complications from mesh removal

Published reports on long-term outcomes of patients after mesh removal surgery are limited. All published studies are retrospective chart or database reviews and substantial heterogeneity exists in terms of both methodology and outcome measures.^{9,41,43,44} Most authors of studies in this area commented on the technical difficulties encountered during mesh excision surgery and the fact that many (in some series, most) patients require two or more surgeries; thus, even in the short term, outcomes are often suboptimal.^{6,9,40–45} Patients who underwent surgery with a primary indication of urethral obstruction had the highest success rates and those whose primary indication was pain had the least successful outcomes. Perioperative decision making is difficult in these patients and is often highly individualized, mostly based on the surgeon's experience and preferences: whether to attempt removal of all the mesh or just the suburethral or vaginal portions and whether to use a synchronous anti-incontinence or urethral reconstruction procedure.⁹ Of note, in patients requiring mesh removal, recurrence of SUI has been reported in 10–60% of individuals.^{9,43,78,84,85,94,100,138} An understanding of the possible outcomes of salvage surgery for mesh complications is critical in enabling accurate decisions regarding informed consent for the use of primary mesh surgery; however, few prospective or registry-based studies with published results currently exist that might address this need.^{9,44,326}

Mesh–body Interactions

Despite the extensive use of polypropylene mesh dating back to the late 1950s,³⁴⁰ in a variety of medical procedures, a paucity of data currently exists regarding the fate of this type of mesh once implanted in humans. Almost all of our knowledge of mesh–body interactions is derived from animal studies; explanted material from patients has been largely neglected as a source of information in this area. After >50 years of use, only a few published studies exist in which investigators actually examined histological changes in mesh explants that had been removed from humans.^{35,341–343}

Despite this general lack of information, in one study of human explanted mesh samples and pathology records from 102 patients, <50% of explanted transvaginal mesh specimens were examined microscopically; however, when microscopy was performed, results of the microscopic examinations usually did not explain the specific complications experienced by the patients.²⁵³ Several studies have confirmed this finding, noting that the assumption that mesh is widely considered to be biologically inert is based on results of short-term animal experiments without corroborating studies in humans.^{342–344} At present, general human tissue interactions with the mesh are known, but we have an incomplete understanding of interactions specific to a mesh material and design as well as the pathophysiology of any complications.

Tissue responses to mesh Inflammatory reactions

The inflammatory response to implanted mesh is non-specific, similar to the foreign-body type of reaction initially described in the late 19th century.^{345,346} However, only since the 1990s have tissue–implant interactions been studied, and, to date, few reports of the mechanisms involved have been published.^{342–344} Immediately after implantation, foreign bodies, including modern implantable polymers, become coated with proteins followed by the appearance of inflammatory cells that migrate into the tissue, owing to the action of inflammatory mediators.^{347,348} As in any tissue injury, the acute phase of inflammation is characterized by the appearance of short-lived neutrophils. Neutrophils are replaced within days by macrophages, which persist indefinitely. The initial phagocyte migration towards the foreign body does not seem to be driven by chemoattractants, but is dependent on the proteins, specifically fibrinogen, coating the implanted objects.^{347,348} The macrophages then either persist and take on an epithelioid appearance or fuse to form multinucleated giant cells. Macrophage fusion occurs in the presence of certain cytokines, when the foreign object is too large to be phagocytosed by a single cell.³⁴⁹ The macrophages are recruited in an attempt to destroy the foreign object and are the main component of the granulomatous inflammation triggered by the foreign body. The macrophages secrete an array of substances, such as bioactive lipids, hydrolytic enzymes, reactive oxygen metabolites and mediators of fibroblast proliferation.^{350,351} In addition to




Figure 5 | Scar encapsulating mesh and surrounding pre-existent normal adipose and muscular tissues. **a** | $\times 2.5$ image of a histological section showing a cross-section of mesh filaments as they appear in section, without colouring. Some filaments were labelled blue by the manufacturer. Adipose tissue had been present in the area before mesh placement. Tissue reaction to surgical injury and the mesh generated scar tissue encapsulating the mesh appears as dense pink collagenous tissue. The scar spans, or bridges, across mesh pores, which is termed bridging fibrosis. In this case a terminal pore contains nonscar adipose tissue (arrow with AT). This section has been labelled with a haematoxylin and eosin stain. **b** | $\times 2.5$ image of a histological section showing cross-sections of mesh filaments. Note that the mesh is surrounded by a halo of fibrous tissue separating it from the pre-existent tissue of the vaginal wall, containing smooth muscle. Smooth muscle is labelled with anti α smooth muscle actin (brown), mesh filaments are filled yellow. The blue colour is a haematoxylin background stain. Abbreviations: AT, adipose tissue; BF, bridging fibrosis.

macrophage-mediated effects, a granulomatous reaction includes T lymphocytes as well as a smaller proportion of B lymphocytes and plasma cells. Each mesh filament ultimately becomes surrounded by a sheath of granulomatous inflammation and the entire mesh structure remains chronically inflamed.^{342,352–354} Three clinically important aspects of mesh-induced inflammation exist: inflammatory mechanisms of pain, stimulation of fibrosis and polypropylene degradation.

Mesh integration

Mesh integration into the tissue is the result of wound repair mechanisms, which aim to restore tissue continuity. In most mammals after the foetal stage of development, the damaged tissue and void spaces are filled with fibrous, or scar tissue. This fibrous tissue functions as a nonspecific universal repair material or filler. In relation to implanted mesh, the body needs to repair tissue that was damaged during surgery, as well as fill the spaces within the mesh structure. The body also needs to repair the tissue damaged by mesh-associated inflammation.

With implanted mesh, granulomatous tissue inhabits the spaces within the mesh structure, such as the pores and interstices between mesh filaments; however, only provided the spaces are large enough to allow tissue

ingrowth.³⁴⁴ During the weeks after SMUS implantation, collagen deposits accumulate, while the fibroblasts acquire contractile filaments and transform into myofibroblasts. The contractile functions of these myofibroblasts together with reduction of extracellular fluid and collagen crosslinking results in wound contraction;^{316,355} the overall aim of wound contraction being to minimize the volume of the maturing scar. In the scar-inhabiting mesh, the contractile forces act on the interlocked mesh–scar composite structure, which results in mesh contraction.^{103,316,356} Maturation of the newly generated fibrous tissue is the next step in the repair process. During this maturation stage, collagen becomes increasingly organized and the density of the microvasculature recedes.³⁵⁵

Initially, the scar is composed of type III collagen, which is replaced by type I collagen as the scar matures. In the transition from type III to type I collagen, the structure is rearranged into cross-linked sheets that run parallel to tension forces.³⁵⁵ The repaired area becomes a hypocellular scar that is then slowly remodeled, which can take ≥ 1 year to complete. Repeated or continuous damage to the tissue can cause the process of repair to be renewed at any stage. Thus, chronic inflammatory conditions can generate a large amount of scar tissue.

With foreign bodies such as mesh, the repair process is complicated by the inflammatory reaction, which is a stimulus for fibrosis. The amount of scar tissue that accumulates is dependent upon counterbalancing processes: stimulation, owing to the presence of a foreign body, and reduction of the scar volume by remodeling. In relation to implanted meshes, fibrous tissue fills the spaces within the mesh structure and surrounds the mesh.^{342,343} The tissue then undergoes contraction and remodeling: the stimulus for fibrosis is, therefore, stronger around the mesh filaments and weaker far from the filaments, that is, in the mesh pores. Some larger pores might include fat or other components of normal connective tissue, while the surrounding filaments are fully encapsulated by the scar (Figure 4).³⁵ Bridging fibrosis can occur in the mesh, where the scar spans or bridges across the pores (Figure 5). Lightweight mesh designs containing pores of several millimetres in diameter have a greater chance of containing normal, nonscar connective tissue in the larger pores of their complex structures.^{35,343} By contrast, heavyweight mesh designs, which are currently used for SMUS devices, lead to the development of a continuous scar plate, which encases all mesh filaments and spans across most of the pores (Figure 5).^{35,343,357} Scar tissue also provides a connection between the composite mesh–scar structure and the surrounding normal tissue.

The process of healing also includes restoration of interrupted innervation and innervation of newly formed tissue. After mesh implantation the processes of reinnervation and/or neoinnervation are not overly affected by the presence of mesh.³⁴ Results of a study published in 2014, investigating samples from patients with inguinal hernia showed that the density of nerve branches in the scar encasing the mesh is similar to that of normal tissue

before surgery and marginally, but not significantly, lower than in the scar formed after non-mesh surgery.³⁴ In addition, nerve branches were observed in the interstices and pores of the mesh, probably growing through the mesh similar to small vessels that were observed to cross the mesh plane.³⁴ This finding indicates that tissues superficial to the mesh might be at least partially dependent on the through-the-mesh neurovascular supply (Figure 6).

Mesh degradation

The authors of several studies have reported degradation of polypropylene in explanted meshes;^{36,341,358–361} however, the question of whether polypropylene degrades *in vivo* has not been fully resolved, despite decades of use.^{37,362} Most conclusions of studies in this area were based on the observations of cracking on the exposed surfaces of explanted mesh filaments, which are usually examined using scanning electron microscopy.^{351–353,355} The explanted tissue examined in these studies was typically fixed in formalin and had to be separated from the mesh using chemical reagents. Alternative hypotheses emerged that the cracking was either of residual biofilms or, if degradation occurred, it was induced by the formalin or cleaning reagents used. However, other studies demonstrated a similar appearance of polypropylene degradation occurring outside of the human body.^{363–365} Before the publication of scanning electron microscopy studies of the mesh surface, authors of an earlier study, published in 1976, assessed the mechanical properties and molecular weight of implanted mesh and concluded that polypropylene degrades *in vivo*.³⁶⁶ In this study, the investigators placed polypropylene implants with and without antioxidant subcutaneously in hamsters to determine the rate of degradation of the implant. They periodically removed specimens during a 5-month test

period and analyzed the samples using infrared spectroscopy and dynamic mechanical testing.³⁶³ The analyses showed that degradation began to occur after only a few days, although several factors suggested that the *in vivo* degradation process was similar to autoxidation that occurs in air or oxygen; in this study, the oxidation process was retarded through the use of an antioxidant.³⁶³ A lack of published studies exists in this area, although the limited evidence suggests that implanted polypropylene undergoes a process of oxidative degradation, in which one of the factors is believed to be oxidative substances generated by macrophages.^{366,367}

Effects of mesh on the tissue

Pain

Scar tissue inhabiting the mesh is not simply an inanimate filler, but a living tissue with its own vascular supply, innervation, fluid and acid–base balance mechanisms and immune response.^{34,253,368} This tissue is subject to pain through normal mechanisms, caused by specific factors: persistent chronic inflammation, nerve ingrowth, tissue compartmentalization within the mesh and nonphysiological attachments to mobile tissues.

Inflammatory mediators cause hypersensitivity to everyday stimuli that leads to pain in response to touch or on movement and, if the stimulus is sufficiently high, can even lead to pain sensations at rest.³⁶⁹ As discussed earlier, implantation of polypropylene meshes invariably results in an inflammatory response, which creates an environment capable of decreasing a patient's pain threshold (Figure 7).

The interlocking and compartmentalized nature is another specific feature of the mesh–scar complex. The ingrown tissue is in a vulnerable position, as it might be subjected to physical compression and distortion within the compartments of mesh pores and folds.³⁴ The risk of




Figure 6 | Tissue interactions with explanted sling materials. **a, b** | $\times 10$ images of a histological section showing a neurovascular bundle penetrating through a mesh pore. This section has been labelled with a haematoxylin and eosin stain. Cross sections of mesh filaments are filled yellow for demonstration purposes. The neurovascular bundle is within a mesh pore, orientated perpendicular to the mesh plane. **c** | $\times 10$ image of a histological section showing muscle interposition between mesh filaments. This section has been anti-desmin-labelled (brown) to highlight the presence of striated muscle. Interlocked striated muscle is commonly observed in explanted transobturator tapes. **d** | $\times 10$ image of a histological section showing α -smooth-muscle-actin-labelled smooth muscle from the vaginal wall, urethra or urinary bladder surrounding the sling material. Depending on the muscle origin, smooth muscle is likely to interact with mesh during physiological contractions (such as those that occur during urination or intercourse). Abbreviations: N, nerve branch; V, blood vessel.




Figure 7 | Inflammatory reaction to the mesh. **a** | $\times 40$ image of a histological section showing a cross-section of mesh filaments surrounded by foreign-body-type inflammation. Epithelioid macrophages (between arrows 'M'), which are the main component of granulomatous inflammation can be observed. A smaller number of lymphocytes ('L') can be seen surrounding the mesh filament. This section has been labelled with a haematoxylin and eosin stain. **b** | $\times 40$ image of a histological section showing a cross-section of mesh filaments, characterized by the presence of neutrophils (multiple neutrophils are scattered in the infiltrate, two are labelled 'N' as examples). Acute inflammation is a feature of bacterial infection and is seen in patients with mesh exposure through the vaginal, urethral or bladder mucosa. Abbreviations: L, lymphocytes; M, macrophages; N, neutrophils.

compression might be a result of external forces, such as moving, bending and penile thrusting during intercourse, as well as a result of increased interstitial fluid pressure within the compartments. We have observed evidence of oedema within pores and deformation pockets (folds) of larger mesh devices indicating fluid imbalance within the mesh compartments (Figure 8). Externally, scar connection to the surrounding tissue might cause distortion and pulling during movement. These forces can act on the entire mesh structure, which in a SMUS has a long course compared with that of a hernia patch. This relatively long length of a SMUS might result in multiple sites of scar attachment to the tissues, which, in the case of TOT slings, includes actively contracting striated muscle.²⁷⁷ These nonphysiological connections are subject to pulling forces that might induce pain, either as a result of muscle contraction or mobility during body movements. Additionally, mesh shrinkage during scar contraction might lead to static tension within and between attached tissues, also contributing to pain.¹⁰³

Many authors have suggested nerve entrapment as a cause of SMUS-related pain. Entrapped nerves have been detected in mesh explants from patients with hernia,^{34,35,370} and nerve branches have also been shown to grow into the mesh interstices in up to 90% of explanted mesh samples from patients with hernia.^{34,35} Nerve entrapment has also been reported in patients fitted with a transvaginal mesh,¹¹² although few published studies exist in this area. In our unreported clinical experience of over 100 explanted mesh specimens, nearly all contained nerve branches of variable calibre (Figure 6). Interestingly, in patients undergoing hernia surgery, prophylactic neurectomy is offered as a method to reduce the incidence of pain after mesh repair.³⁷¹

Some patients fitted with a SMUS report pain that is associated with specific movements or activities. The observation of interlocking of the mesh and striated muscle—resulting in muscle contraction and traction on entrapped nerves—offers a plausible hypothesis to explain this phenomenon in patients fitted with TOT slings. We have also observed interposition of smooth muscle, which might contribute to dyspareunia as the vaginal walls contract during intercourse (Figure 6).

Dyspareunia

Direct pressure and a wide range of tissue movement during sexual intercourse both pose additional risks to patients with a SMUS. The vaginal mucosa is one of the most densely innervated parts of the human body. Thus, if this mucosa overlies a stiffened mesh–scar structure, the nerve branches and receptors are subject to compression against the stiffened SMUS during intercourse. In addition, tissue movement on either side of the scar plate can cause traction and distortion of the mesh–scar structure. Findings of a study published in 2010³⁸ demonstrated the existence of a new complication unique to patients fitted with TOT slings, termed banding, which is a palpable firm scar in the para-urethral folds that was associated with dyspareunia in four of 12 sexually active women who were found to have banding on examination. Unfortunately, no data were presented describing the histopathology of excised tissue that was removed owing to painful banding, although little imagination is required to understand how this effect could cause pain (Figure 9).

Mesh exposure in the vaginal wall

Mesh erosion through the vaginal mucosa can result in a large variety of tissue responses, ranging from no detectable changes to substantial acute inflammation (Figure 7) and even formation of small abscesses. The dense acute inflammation almost always signifies bacterial infection.³⁷² From our unpublished experience, the site of mesh exposure also has a variable amount of granulated tissue. These changes correlate with the presenting symptoms, which range from no complaints to vaginal discharge, bleeding, dyspareunia and feeling of the exposed edge of the mesh by the sexual partner.

The mechanisms of mesh exposure and/or extrusion through the vaginal wall have not been well studied; however, an approximately 26-fold increase in vaginal and bladder exposure, extrusion or erosion when there had been a vaginal or bladder trocar perforation during the original surgery is known to exist.³¹ Other risk factors have also been identified including patients' having undergone prior vaginal surgeries, larger incisions, smoking, diabetes mellitus of either type, pelvic exposure to radiation and older patient age.^{96,124,321} The existence of these risk factors suggests that poor healing, lowered antibacterial immunity, insufficient vascularization, and scarring are all possible causes contributing to mesh exposure. In terms of mesh-specific factors, solid silicone strips or silicone-coated meshes have higher rates of erosion, indicating that choice of material

can affect the risk of vaginal mesh exposure.^{105,373,374} Compared with other SMUS materials, silicone has limited adhesion to the tissues, which possibly enables more movement of the mesh, or tissue detachment. Patients fitted with a SMUS with a design that incorporates microporous materials with low tissue adherence, such as Gore-Tex® also have higher rates of mesh exposure through the vaginal wall,³⁷⁵ compared with those of patients fitted with a SMUS that incorporates polypropylene mesh, which has larger pores.^{301,356,376} In addition to the lower tissue adherence that enables mesh movement within the tissue, meshes with smaller pores do not allow tissue growth through the mesh. This lack of tissue growth likely interferes with vascularization and innervation of the overlying mucosa, which might lead to dystrophic changes and poor resistance to infection and necrosis.

Results of experiments conducted in animal models showed that the rate of mesh erosion was also dependent on the size of mesh implant, with animals implanted with larger mesh patches having a higher risk of exposure.^{356,376} The higher exposure rate of larger mesh implants was likely a result of higher risks of mesh migration. Deformation was associated with the use of larger patches, more interference with vascularization and innervation of the overlying mucosa and the presence of larger volume of inflammation and/or fibrosis. Implantation with a smaller area of mesh might result in less risk of exposure, assuming that exposure is an entirely random event. In our unreported clinical experience of over 100 explanted polypropylene slings, we frequently observed that the exposed part of the explanted mesh was a curled edge piercing through the mucosa, suggesting that edge curling is also a mechanism of vaginal mesh exposure. In addition to the internal properties of a knitted structure, outward pressure of the tissue can act to curl the edge. Interestingly, resection of an exposed part of the mesh, either an edge or a mid-portion, leaves new edges that can also curl and become exposed. In our experience many mesh exposures, SMUS-related or implant-related pelvic organ prolapses recur after trimming of the exposed part.

Mesh migration

Mesh exposure through the vaginal wall seems to have several potential causes. Mesh erosion through the urethral or bladder mucosa reflects mesh migration (or incorrect SMUS placement). Mesh migration through the tissues and into the adjacent organs has been described when used in patients with hernia, in which two types of mesh migration have been suggested to occur: primary migration of unsecured mesh towards areas of least tissue resistance and secondary migration through transanatomical planes. The latter is facilitated by tissue forces acting to displace mesh while remodelling-induced and inflammation-induced tissue resorption enable this movement.^{377,378} In patients with SMUS, mesh migration typically occurs into or through the urethral wall, which is a secondary type of migration.^{9,10,39,41,326} Excessive tensioning of the sling can act to




Figure 8 | Oedema within mesh compartments. ×20 image of a histological section showing oedema in mesh compartments, note separation of collagen and low density of tissue in the area of the oedema (E). Oedema is usually seen in semi-enclosed mesh compartments. Mesh filaments are filled yellow in this image. This section has been labelled with a haematoxylin and eosin stain. Abbreviation: E, oedema

displace the mesh into the urethra, whereas an inflammatory reaction to a foreign body and the general ability of tissues to remodel under chronic pressures can enable mesh migration. Remnants of partially excised SMUS can potentially migrate in directions other than into the urethra.

Mesh deformation

Mesh deformation, in which a part of mesh moves from its original or intended position, is related to mesh migration. In an *in vivo* study using white rabbits, Amid type I (Marlex®) meshes were found to be more likely to fold or curl at the edges in comparison to Amid type II meshes (Teflon®).³⁷⁹ Similar to mesh migration, deformation can be primary, as a result of intraoperative or perioperative folding and edge curling of an unsecured mesh, or secondary, occurring after tissue ingrowth. Secondary wrinkling and folding of the mesh is attributed largely to mesh–scar contraction.³¹⁶ For transvaginal applications, folding and bunching of the mesh is frequently observed in patients with pelvic organ prolapse, who are often fitted with large devices, whereas mesh deformation of SMUS devices is typically limited to edge curling (Figure 9).³⁸⁰ Edge curling of knitted mesh materials has been noticed following their surgical use in patients with hernia, where the edges can be secured by stitching; however, transvaginal devices have all edges unsecured. Narrow sling tapes might also show signs of fraying and curling of the edges when stretched. To address this risk of sling deformation, SMUS manufacturers have used heat treatment approaches, with variable degree of success.³⁸¹ As we note, rotation of a frayed edge towards the mucosa is another possible




Figure 9 | Curling of the edges of explanted sling materials. **a** | Segment of a sling, which was explanted with very little adherent tissue and a structure that is readily visible. **b** | Segment of a mesh sling, which was excised with adherent tissue remaining attached to the sling material.

mechanism by which mucosal exposure of the mesh might take place.

Mesh stiffening

Elasticity and flexibility of knitted meshes is dependent on bending and movement of the mesh filaments. The extent of freedom of movement is often substantially reduced by the ingrowth of collagenous scar tissue. At the same time, the embedded mesh acts to reinforce the scar tissue, thus limiting native flexibility of the scar.^{380,382–384} The resultant mesh–scar composite structure is stiffer than the original new mesh. The extent of the resultant increase in stiffness is dependent on mesh design, including the physical characteristics of the material and the amount of induced fibrous scar.^{294,316,382–384} For SMUS devices, excessive tightening and connection to the surrounding tissues might limit mobility and add to the structural stiffness. This phenomenon has been observed in the clinic, where it is referred to as ‘banding’.³⁸ Mesh stiffening is likely to occur owing to degradation of polypropylene, as the degraded layer often shows embrittlement.^{357–359,361,366} This component of stiffening is expected to increase over time.

Urinary symptoms

SMUS are designed to support the urethra; however, as discussed previously, the amount of pressure can become excessive owing to mesh contraction and reduction of the area of support. A stretched mesh has a reduced width, which, together with edge curling has been described as ‘roping’ (Figure 9).³⁸⁵ A stiffened, over-tightened sling, therefore, has limited elasticity and cannot accommodate a full range of changes in tension. SMUS tension changes dynamically during cough, sexual intercourse and other physiological processes, which adds to the static pressure on the urethra. The result is urinary retention and transmigration of the mesh into, or through the urethral wall. Interestingly, mesh removal does not necessarily lead to recurrent SUI.^{369,370} This finding suggests that scarring around the mesh, which remains after mesh explantation, is sufficient to maintain continence in some patients.

Polypropylene degradation products

The breakdown of mesh is expected to result in the presence of small molecular complexes and chemical products of degradation, as is the case for any polymerized hydrocarbon. *In vitro* thermal degradation of polypropylene at high temperatures produces an array of organic molecules such as acids, ketones, ethers, aldehydes, alcohols and smaller hydrocarbons.³⁸⁶ The *in vitro* conditions required for thermal degradation, however, are different to those observed under *in vivo* conditions, and we are not aware of any studies that either simulated body conditions or conducted chemical analysis of explanted tissue. An assumption can be made that, to some extent, any combination of the degradation products detected during thermal or other types of degradation can be produced in the tissue. Additionally, additives used to stabilize the polymer might theoretically leach into the surrounding tissue.

Accumulations of polypropylene degradation products are expected to be confined within the scar capsule and have more local, rather than systemic, effects on the body, owing to their fibrous encapsulation; however, no published studies currently address this point. The degradation products might act as an additional stimulus for the chronic inflammatory response. Accumulation and toxicity of these degradation products might cause tissue damage and contribute to the continuous remodelling around the mesh filaments and extension of fibrosis.³⁸⁷

Tumorigenicity

Three cases of cancer that might be associated with implanted polypropylene mesh have been reported in humans. Two patients had squamous cell cancers 6 years and 22 years after mesh hernia repairs, respectively;³⁸⁸ in addition, an inflammatory myofibroblastic tumour following implantation of an RP sling has also been reported.³⁸⁴ In the patients who had mesh hernia repairs, both had a complicated clinical course involving chronic mesh exposure and infection.³⁸³ Chronic skin wounds are an established risk factor for squamous cell carcinoma. The one known patient with an RP-sling-related tumour had a myofibroblastic neoplasm with local recurrence potential, which is considered an intermediate state between a benign tumour and sarcoma. Potential risks of tumorigenesis in patients with SMUS include chronic mucosal erosions, chronic inflammation surrounding the mesh and the possible presence of degradation products and polypropylene additives released into the tissue. Mutagenic effects, in general can take many years to accumulate and then a long period to cause a neoplasm. Detecting any oncogenic effects of SMUS implants would require a large cohort of patients with the same type of implant, and these patients would have to be followed up for a sufficiently long period of time, most likely at least 15 years. The long-term use of hernia meshes has not revealed a significant oncogenic risk; however, the constant introduction of new mesh designs further complicates investigations of mesh-related cancer risks, as these new

designs probably vary in terms of the chemical composition of the polypropylene used. In general, based on our knowledge of tumorigenesis, three tissue types might be affected by introduction of a SMUS: epithelium; soft tissue; and lymphocytes, which could result in malignant transformation into carcinoma, sarcoma and lymphoma, respectively.

As described, a small risk of developing squamous cell carcinoma associated with chronic mesh exposure has been reported in patients with hernia meshes.³⁸⁸ In transvaginal applications of similar materials, chronic erosions might, therefore, increase the risk of squamous cell carcinoma. The importance of concurrent local infections with high-risk variants of human papillomavirus needs to be studied, as these might have a synergistic effect in increasing a patient's cancer risk.

Carcinogenic effects of polypropylene, specifically leading to the development of sarcomas, have been studied in animal models. In rodents, implantation with flat polypropylene plates resulted in higher tumorigenicity than placement of porous materials.³⁸⁹ A study of implanted polypropylene meshes in mice concluded that the risk of carcinogenesis following mesh implantation, if existent, is not immediate; however the follow-up duration of this study was only 2 years.³⁹⁰ Another group of researchers implanted transponders made of polypropylene into carcinogen-sensitive *p53*⁺ transgenic mice and observed development of sarcomas in 10% of animals within 6 months of exposure.³⁹¹ Other reports have corroborated these findings.^{392–394} The basic research and clinical data suggest that implantation of polypropylene mesh might increase the risk of sarcoma, however if a risk is present in humans it is likely to be very low.

A potential risk of lymphoma needs to be considered in patients with any prosthetic implants, including SMUS, as this effect has been well documented in women with breast implants.^{395,396} The exact aetiology of lymphoma owing to breast implants is not presently known, and this increased risk was detected in association with either saline or silicone implants. The increased risk of lymphoma might be related to an inflammatory reaction to the implants, rather than to the material of the devices; therefore, this risk might also be relevant to a large range of other implants that induce inflammatory responses, including SMUS. The large size of breast implants relative to most other implanted materials and the high volume of use for over 30 years might explain why this small, specific risk became detectable. Whether or not the same risk exists in patients with SMUS is currently unknown. In women with breast implants, the average time between implantation and a diagnosis of lymphoma is reported to be 9 years (range 1–32 years).³⁹⁶

Conclusion

In the words of the astronomer Carl Sagan—"The absence of evidence is not evidence of absence".³⁹⁷ With respect to the safety of sling surgery, the lack of good studies about the incidence and severity of SMUS

complications is not evidence that these complications are uncommon, nor is it evidence that they are not serious. The effectiveness of synthetic slings remains unchallenged, although, as this Review documents, an increasing body of evidence exists that serious and sometimes lifestyle-altering complications are under-reported and underappreciated by doctors and patients alike. The true incidence of SMUS-related complications is unknown, owing, in no small part, to the poor overall quality of the studies. Nevertheless, we have calculated the minimum risks: revision surgery for erosion and obstruction alone, 5.6%; chronic pain, 4.3%; recurrent or persistent SUI, 5.3%; and *de novo* refractory OAB 3.9% (Box 1). These data are not mutually exclusive, although we calculated the overall risk of a serious complication or surgical failure to be 12.5%. We emphasize, though, that these data represent the absolute minimum rate of complications reported in the literature; the actual rate might be considerably higher.

Urologists can and must do better in assessing the long-term safety of SMUS surgery and in developing better methods of monitoring patients and assessing the outcomes of treatment for complications, so that both patients and physicians can be advised of the true risks associated with a SMUS.

Review criteria

A systematic review of the English language literature was performed in August 2014 to investigate the published efficacy, effectiveness and complications of SMUS. The search used a complex search strategy of the Medline database, including medical subject heading (MeSH) and free-text protocols. The MeSH search combined the terms "mid urethral sling", "midurethral sling", "suburethral sling", "urethral sling", "mid urethral slings", "midurethral slings", "suburethral slings", "urethral slings" and "follow-up study". Multiple free-text searches included the terms "Urinar*incont*", "TVT", "tension-free vaginal tape*", "Tension-free vaginal sling*", "Transobturator tape*", "Transobturator sling*", "TVT-obturator", "TVT-O", "TVT secure", "miniarc", "abbrevio", "TOT", "suprapubic arc sling*", "SPARC sling*", "intravaginal slingplasty", "IVS sling", "Raz sling", "Uratape", "ObTAPE", "Prepubic sling*", "Prepubic TVT", "Prepubic tape*", "PelviLace", "Ureter", "Aris", "In-Fast", "Monarc", "I-Stop", "urethral reconstruction", "urethrovaginal fistula", "Obtape", "gortex sling", "silastic sling", "mersilene sling", "marlex sling", "vesicovaginal fistula", "BioArc" individually in the fields title and abstract of the records. Subsequently, the search was limited to only human patients. A total of 995 records were retrieved from Medline, 249 were included. Six of the authors reviewed the full texts to select relevant papers. Discrepancies were solved by open discussion. Once the citations were accrued and the papers read, the bibliographies were cross-checked for any relevant citations that were missed in the initial search, which totalled an additional 88 articles. Only articles published since 2007 were included in the reporting of complications to update and expand upon a review published in 2008.³³⁴ For the section on mesh-body interactions, the search was not limited to humans nor was there a limit on publication date.

1. Dmochowski, R. R. *et al.* Update of AUA guideline on the surgical management of female stress urinary incontinence. *J. Urol.* **183**, 1906–1914 (2010).
2. Blaivas, J. G. & Jacobs, B. Z. Pubovaginal fascial sling for the treatment of complicated stress urinary incontinence. *J. Urol.* **145**, 1214–1218 (1991).
3. Appell, R. A. Primary slings for everyone with genuine stress incontinence? The argument for... *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **9**, 249–251 (1998).
4. Chaikin, D. C., Rosenthal, J. & Blaivas, J. G. Pubovaginal fascial sling for all types of stress urinary incontinence: long-term analysis. *J. Urol.* **160**, 1312–1316 (1998).
5. Nager, C., Tulikangas, P., Miller, D., Rovner, E. & Goldman, H. Position statement on mesh midurethral slings for stress urinary incontinence. *Female Pelvic Med. Reconstr. Surg.* **20**, 123–125 (2014).
6. Jonsson Funk, M., Siddiqui, N. Y., Pate, V., Amundsen, C. L. & Wu, J. M. Sling revision/removal for mesh erosion and urinary retention: long-term risk and predictors. *Am. J. Obstet. Gynecol.* **208**, 73.e1–e7 (2013).
7. Chughtai, B. I. *et al.* Midurethral sling is the dominant procedure for female stress urinary incontinence: analysis of case logs from certifying American urologists. *Urology* **82**, 1267–1271 (2013).
8. Ogah, J., Cody, D. J. & Rogerson, L. Minimally invasive synthetic suburethral sling operations for stress urinary incontinence in women: a short version Cochrane review. *Neurourol. Urodyn.* **30**, 284–291 (2011).
9. Blaivas, J. *et al.* Salvage surgery after failed treatment of synthetic mesh sling complications. *J. Urol.* **190**, 1281–1286 (2013).
10. Deng, D., Rutman, M., Raz, S. & Rodriguez, L. Presentation and management of major complications of midurethral slings: are complications under-reported? *Neurourol. Urodyn.* **26**, 46–52 (2007).
11. Ulmsten, U. & Petros, P. Intravaginal slingplasty (IVS): an ambulatory surgical procedure for treatment of female urinary incontinence. *Scand. J. Urol. Nephrol.* **29**, 75–82 (1995).
12. Delorme, E. Transobturator urethral suspension: mini-invasive procedure in the treatment of stress urinary incontinence in women [French]. *Prog. Urol.* **11**, 1306–1313 (2001).
13. Latthe, P., Foon, R. & Toozs-Hobson, P. Transobturator and retropubic tape procedures in stress urinary incontinence: a systematic review and meta-analysis of effectiveness and complications. *BJOG* **114**, 522–531 (2007).
14. Latthe, P. M., Singh, P., Foon, R. & Toozs-Hobson, P. Two routes of transobturator tape procedures in stress urinary incontinence: a meta-analysis with direct and indirect comparison of randomized trials. *BJU Int.* **106**, 68–76 (2010).
15. Novara, G. *et al.* Updated systematic review and meta-analysis of the comparative data on colposuspensions, pubovaginal slings, and midurethral tapes in the surgical treatment of female stress urinary incontinence. *Eur. Urol.* **58**, 218–238 (2010).
16. Nilsson, C., Palva, K., Aarnio, R., Morcos, E. & Falconer, C. Seventeen years' follow-up of the tension-free vaginal tape procedure for female stress urinary incontinence. *Int. Urogynecol. J.* **24**, 1265–1269 (2013).
17. Cheng, D. & Liu, C. Tension-free vaginal tape-obturator in the treatment of stress urinary incontinence: a prospective study with five-year follow-up. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **161**, 228–231 (2012).
18. Serati, M. *et al.* TVT-O for the treatment of pure urodynamic stress incontinence: efficacy, adverse effects, and prognostic factors at 5-year follow-up. *Eur. Urol.* **63**, 872–878 (2013).
19. Li, B., Zhu, L., Lang, J., Fan, R. & Xu, T. Long-term outcomes of the tension-free vaginal tape procedure for female stress urinary incontinence: 7-year follow-up in China. *J. Minim. Invasive Gynecol.* **19**, 201–205 (2012).
20. Angioli, R. *et al.* Tension-free vaginal tape versus transobturator suburethral tape: five-year follow-up results of a prospective, randomised trial. *Eur. Urol.* **58**, 671–677 (2010).
21. Olsson, I., Abrahamsson, A. K. & Kroon, U. B. Long-term efficacy of the tension-free vaginal tape procedure for the treatment of urinary incontinence: a retrospective follow-up 11.5 years post-operatively. *Int. Urogynecol. J.* **21**, 679–683 (2010).
22. Svenningsen, R., Staff, A. C., Schiotz, H. A., Western, K. & Kulseng-Hanssen, S. Long-term follow-up of the retropubic tension-free vaginal tape procedure. *Int. Urogynecol. J.* **24**, 1271–1278 (2013).
23. Groutz, A. *et al.* Ten-year subjective outcome results of the retropubic tension-free vaginal tape for treatment of stress urinary incontinence. *J. Minim. Invasive Gynecol.* **18**, 726–729 (2011).
24. Shumaker, S. A., Wyman, J. F., Uebersax, J. S., McClish, D. & Fantl, J. A. Health-related quality of life measures for women with urinary incontinence: the Incontinence Impact Questionnaire and the Urogenital Distress Inventory. Continence Program in Women (CPW) Research Group. *Qual. Life Res.* **3**, 291–306 (1994).
25. Blaivas, J., Chancellor, M. B., Verhaaren, M. & Weiss, J. P. Atlas of Urodynamics, 2nd edn Ch. 5, 46–55 (Wiley-Blackwell, 2007).
26. Athanasiou, S. *et al.* Seven years of objective and subjective outcomes of transobturator (TVT-O) vaginal tape: why do tapes fail? *Int. Urogynecol. J.* **25**, 219–225 (2014).
27. Groutz, A., Rosen, G., Gold, R., Lessing, J. B. & Gordon, D. Long-term outcome of transobturator tension-free vaginal tape: efficacy and risk factors for surgical failure. *J. Womens Health (Larchmt)* **20**, 1525–1528 (2011).
28. Groutz, A. *et al.* Noninvasive outcome measures of urinary incontinence and lower urinary tract symptoms: a multicentre study of micturition diary and pad tests. *J. Urol.* **164**, 698–701 (2000).
29. Ankardal, M., Heiwall, B., Lausten-Thomsen, N., Carnelid, J. & Milsom, I. Short- and long-term results of the tension-free vaginal tape procedure in the treatment of female urinary incontinence. *Acta Obstet. Gynecol. Scand.* **85**, 986–992 (2006).
30. Albo, M. *et al.* Treatment success of retropubic and transobturator mid urethral slings at 24 months. *J. Urol.* **188**, 2281–2287 (2012).
31. Osborn, D. J. *et al.* Analysis of patient and technical factors associated with midurethral sling mesh exposure and perforation. *Int. J. Urol.* **11**, 1167–1170 (2014).
32. Culligan, P. *et al.* Bacterial colony counts during vaginal surgery. *Infect. Dis. Obstet. Gynecol.* **11**, 161–165 (2003).
33. Wang, A., Lee, L., Lin, C. & Chen, J. A histologic and immunohistochemical analysis of defective vaginal healing after continence taping procedures: a prospective case-controlled pilot study. *Am. J. Obstet. Gynecol.* **191**, 1868–1874 (2004).
34. Bendavid, R., Lou, W., Koch, A. & Iakovlev, V. Mesh-related SIN syndrome. A surreptitious irreversible neuralgia and its morphologic background in the etiology of post-herniorrhaphy pain. *Int. J. Clin. Med.* **5**, 799–810 (2014).
35. Klosterhalfen, B., Junge, K. & Klinge, U. The lightweight and large porous mesh concept for hernia repair. *Expert Rev. Med. Devices* **2**, 103–117 (2005).
36. Ostergard, D. Degradation, infection and heat effects on polypropylene mesh for pelvic implantation: what was known and when it was known. *Int. Urogynecol. J.* **22**, 771–774 (2011).
37. Sternschuss, G., Ostergard, D. & Patel, H. Post-implantation alterations of polypropylene in the human. *J. Urol.* **188**, 27–32 (2012).
38. Chohan, H. J., Hutchings, T. B. & Rooney, K. E. Dyspareunia associated with paraurethral banding in the transobturator sling. *Am. J. Obstet. Gynecol.* **202**, 481.e1–e5 (2010).
39. Brubaker, L. *et al.* Adverse events over two years after retropubic or transobturator midurethral sling surgery: findings from the trial of midurethral slings (TOMUS) study. *Am. J. Obstet. Gynecol.* **205**, 498.e1–e6 (2011).
40. Dunn, G. E. *et al.* Changed women: the long-term impact of vaginal mesh complications. *Female Pelvic Med. Reconstr. Surg.* **20**, 131–136 (2014).
41. Blaivas, J. & Mekel, G. Management of urinary fistulas due to midurethral sling surgery. *J. Urol.* **192**, 1137–1142 (2014).
42. Shah, H. N. & Badlani, G. H. Mesh complications in female pelvic floor reconstructive surgery and their management: A systematic review. *Indian J. Urol.* **28**, 129–153 (2012).
43. Shah, K., Nikolavsky, D., Gilsdorf, D. & Flynn, B. J. Surgical management of lower urinary mesh perforation after mid-urethral polypropylene mesh sling: mesh excision, urinary tract reconstruction and concomitant pubovaginal sling with autologous rectus fascia. *Int. Urogynecol. J.* **24**, 2111–2117 (2013).
44. Abbott, S. *et al.* Evaluation and management of complications from synthetic mesh after pelvic reconstructive surgery: a multicentre study. *Am. J. Obstet. Gynecol.* **210**, 163.e1–e8 (2014).
45. Stanford, E. & Paraiso, M. A comprehensive review of suburethral sling procedure complications. *J. Minim. Invasive Gynecol.* **15**, 132–145 (2008).
46. Aigmueiller, T. *et al.* Ten-year follow-up after the tension-free vaginal tape procedure. *Am. J. Obstet. Gynecol.* **205**, 496.e1–e5 (2011).
47. Ala-Nissila, S., Haarala, M. & Makinen, J. Tension-free vaginal tape - a suitable procedure for patients with recurrent stress urinary incontinence. *Acta Obstet. Gynecol. Scand.* **89**, 210–216 (2010).
48. Al-Singary, W. *et al.* Trans-obturator tape for incontinence: a 3-year follow-up. *Urol. Int.* **78**, 198–201 (2007).
49. Ammendrup, A. C., Jorgensen, A., Sander, P., Ottesen, B. & Lose, G. A Danish national survey of women operated with mid-urethral slings in 2001. *Acta Obstet. Gynecol. Scand.* **88**, 1227–1233 (2009).
50. Andonian, S., St-Denis, B., Lemieux, M. C. & Corcos, J. Prospective clinical trial comparing Obtape and DUPS to TVT: one-year safety and efficacy results. *Eur. Urol.* **52**, 245–251 (2007).
51. Anger, J. T., Rodriguez, L. V., Wang, Q., Pashos, C. L. & Litwin, M. S. The role of preoperative testing on outcomes after sling surgery for stress urinary incontinence. *J. Urol.* **178**, 1364–1368 (2007).

52. Araco, F. *et al.* Results 1 year after the Reemex system was applied for the treatment of stress urinary incontinence caused by intrinsic sphincter deficiency. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 783–786 (2008).
53. Ark, C., Gungorduk, K. & Celebi, I. Results of TOT operations alone and combined with other vaginal surgical procedures. *Aust. N. Z. J. Obstet. Gynaecol.* **50**, 550–555 (2010).
54. Ballert, K. N., Kanofsky, J. A. & Nitti, V. W. Effect of tension-free vaginal tape and TVT-obturator on lower urinary tract symptoms other than stress urinary incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 335–340 (2008).
55. Barber, M. D. *et al.* Transobturator tape compared with tension-free vaginal tape for the treatment of stress urinary incontinence: a randomized controlled trial. *Obstet. Gynecol.* **111**, 611–621 (2008).
56. Barber, M. D. *et al.* Single-incision mini-sling compared with tension-free vaginal tape for the treatment of stress urinary incontinence: a randomized controlled trial. *Obstet. Gynecol.* **119**, 328–337 (2012).
57. Biggs, G. Y., Ballert, K. N., Rosenblum, N. & Nitti, V. Patient-reported outcomes for tension-free vaginal tape-obturator in women treated with a previous anti-incontinence procedure. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 331–335 (2009).
58. Chae, H. D. *et al.* A comparative study of inside-in and outside-out transobturator tape procedures for stress urinary incontinence. *Gynecol. Obstet. Invest.* **70**, 200–205 (2010).
59. Charalambous, S. *et al.* Transvaginal vs transobturator approach for synthetic sling placement in patients with stress urinary incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 357–360 (2008).
60. Chen, H. Y., Ho, M., Hung, Y. C. & Huang, L. C. Analysis of risk factors associated with vaginal erosion after synthetic sling procedures for stress urinary incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 117–121 (2008).
61. Cho, M. K. *et al.* Comparison of the clinical and quality-of-life outcomes after the inside-out TVT-O procedure with or without concomitant transvaginal gynaecological surgery. *J. Obstet. Gynaecol.* **32**, 280–284 (2012).
62. Collinet, P. *et al.* The safety of the inside-out transobturator approach for transvaginal tape (TVT-O) treatment in stress urinary incontinence: French registry data on 984 women. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 711–715 (2008).
63. Costantini, E. *et al.* Preoperative Valsalva leak point pressure may not predict outcome of mid-urethral slings. Analysis from a randomized controlled trial of retropubic versus transobturator mid-urethral slings. *Int. Braz. J. Urol.* **34**, 73–81 (2008).
64. Deffieux, X. *et al.* Transobturator TVT-O versus retropubic TVT: results of a multicentre randomized controlled trial at 24 months follow-up. *Int. Urogynecol. J.* **21**, 1337–1345 (2010).
65. Dobson, A. *et al.* Trans-obturator surgery for stress urinary incontinence: 1-year follow-up of a cohort of 52 women. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 27–32 (2007).
66. Feng, C. L., Chin, H. Y. & Wang, K. H. Transobturator vaginal tape inside out procedure for stress urinary incontinence: results of 102 patients. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 1423–1427 (2008).
67. Giberti, C., Gallo, F., Cortese, P. & Schenone, M. Transobturator tape for treatment of female stress urinary incontinence: objective and subjective results after a mean follow-up of two years. *Urology* **69**, 703–707 (2007).
68. Heinonen, P., Ala-Nissila, S., Kiilholma, P. & Laurikainen, E. Tension-free vaginal tape procedure without preoperative urodynamic examination: long-term outcome. *Int. J. Urol.* **19**, 1003–1009 (2012).
69. Houwert, R. M., Renes-Zijl, C., Vos, M. C. & Vervest, H. A. TVT-O versus Monarc after a 2–4-year follow-up: a prospective comparative study. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 1327–1333 (2009).
70. Kaelin-Gambirasio, I., Jacob, S., Boulvain, M., Dubuisson, J. B. & Dallenbach, P. Complications associated with transobturator sling procedures: analysis of 233 consecutive cases with a 27 months follow-up. *BMC Womens Health* **9**, 28 (2009).
71. Killingsworth, L. B. *et al.* One-year outcomes of tension-free vaginal tape (TVT) mid-urethral slings in overweight and obese women. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 1103–1108 (2009).
72. Kocjancic, E. *et al.* Transobturator tape in the management of female stress incontinence: clinical outcomes at medium term follow-up. *Urol. Int.* **80**, 275–278 (2008).
73. Lee, J. H. *et al.* Modified transobturator tape (canal transobturator tape) surgery for female stress urinary incontinence. *J. Urol.* **181**, 2616–2621 (2009).
74. Liapis, A., Bakas, P. & Creatsas, G. Tension-free vaginal tape in the management of recurrent urodynamic stress incontinence after previous failed midurethral tape. *Eur. Urol.* **55**, 1450–1455 (2009).
75. Liapis, A., Bakas, P. & Creatsas, G. Long-term efficacy of tension-free vaginal tape in the management of stress urinary incontinence in women: efficacy at 5- and 7-year follow-up. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 1509–1512 (2008).
76. Lim, J., Cornish, A. & Carey, M. P. Clinical and quality-of-life outcomes in women treated by the TVT-O procedure. *BJOG* **113**, 1315–1320 (2006).
77. Meschia, M., Pifarotti, P., Gattei, U. & Bertozzi, R. Tension-free vaginal tape: analysis of risk factors for failures. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 419–422 (2007).
78. Misrai, V. *et al.* Surgical resection for suburethral sling complications after treatment for stress urinary incontinence. *J. Urol.* **181**, 2198–2202 (2009).
79. Mostafa, A. *et al.* Multicentre prospective randomized study of single-incision mini-sling vs tension-free vaginal tape-obturator in management of female stress urinary incontinence: a minimum of 1-year follow-up. *Urology* **82**, 552–559 (2013).
80. Mostafa, A., Madhuvrata, P. & Abdel-Fattah, M. Preoperative urodynamic predictors of short-term voiding dysfunction following a transobturator tension-free vaginal tape procedure. *Int. J. Gynaecol. Obstet.* **115**, 49–52 (2011).
81. Nazemi, T. M. *et al.* Minimum 24-month followup of the sling for the treatment of stress urinary incontinence. *J. Urol.* **179**, 596–599 (2008).
82. Neuman, M., Sosnovski, V., Goralnik, S., Diker, B. & Bornstein, J. Comparison of two inside-out transobturator suburethral sling techniques for stress incontinence: early postoperative thigh pain and 3-year outcomes. *Int. J. Urol.* **19**, 1103–1107 (2012).
83. Novi, J. M. & Mulvihill, B. H. Surgical intervention for stress urinary incontinence: comparison of midurethral sling procedures. *J. Am. Osteopath. Assoc.* **108**, 634–638 (2008).
84. Padmanabhan, P. *et al.* Approach to management of iatrogenic foreign bodies of the lower urinary tract following reconstructive pelvic surgery. *J. Urol.* **187**, 1685–1690 (2012).
85. Pikaart, D. P., Miklos, J. R. & Moore, R. D. Laparoscopic removal of pubovaginal polypropylene tension-free tape slings. *JSLs* **10**, 220–225 (2006).
86. Porena, M. *et al.* Tension-free vaginal tape versus transobturator tape as surgery for stress urinary incontinence: results of a multicentre randomised trial. *Eur. Urol.* **52**, 1481–1490 (2007).
87. Pushkar, D. Y., Godunov, B. N., Gvozdev, M. & Kasyan, G. R. Complications of mid-urethral slings for treatment of stress urinary incontinence. *Int. J. Gynaecol. Obstet.* **113**, 54–57 (2011).
88. Reich, A., Wiesner, K., Kohorst, F., Kreienberg, R. & Flock, F. Comparison of transobturator vaginal tape and retropubic tension-free vaginal tape: clinical outcome and sonographic results of a case-control study. *Gynecol. Obstet. Invest.* **68**, 137–144 (2009).
89. Ross, S. *et al.* Transobturator tape compared with tension-free vaginal tape for stress incontinence: a randomized controlled trial. *Obstet. Gynecol.* **114**, 1287–1294 (2009).
90. Scheiner, D. A. *et al.* Twelve months effect on voiding function of retropubic compared with outside-in and inside-out transobturator midurethral slings. *Int. Urogynecol. J.* **23**, 197–206 (2012).
91. Schierlitz, L. *et al.* Three-year follow-up of tension-free vaginal tape compared with transobturator tape in women with stress urinary incontinence and intrinsic sphincter deficiency. *Obstet. Gynecol.* **119**, 321–327 (2012).
92. Siddiqui, K., Raj, H., Flynn, R. J., Grainger, R. & Thornhill, J. A. Minimally invasive treatment of female stress urinary incontinence: 100 cases using SPARC sling. *Ir. J. Med. Sci.* **177**, 39–42 (2008).
93. Sung, V. W., Kassir, N. & Raker, C. A. Improvements in physical activity and functioning after undergoing midurethral sling procedure for urinary incontinence. *Obstet. Gynecol.* **120**, 573–580 (2012).
94. Sweat, S. D. *et al.* Polypropylene mesh tape for stress urinary incontinence: complications of urethral erosion and outlet obstruction. *J. Urol.* **168**, 144–146 (2002).
95. Ugurlucan, F. G., Erkan, H. A., Onal, M. & Yalcin, O. Randomized trial of graft materials in transobturator tape operation: biological versus synthetic. *Int. Urogynecol. J.* **24**, 1315–1323 (2013).
96. Van Baelen, A. A. & Delaere, K. P. Repeat transobturator tape after failed mid-urethral sling procedure: follow-up with questionnaire-based assessment. *Urol. Int.* **83**, 399–403 (2009).
97. Verbrugghe, A., De Ridder, D. & Van der Aa, F. A repeat mid-urethral sling as valuable treatment for persistent or recurrent stress urinary incontinence. *Int. Urogynecol. J.* **24**, 999–1004 (2013).
98. Waltregny, D. *et al.* TVT-O for the treatment of female stress urinary incontinence: results of a prospective study after a 3-year minimum follow-up. *Eur. Urol.* **53**, 401–408 (2008).
99. Yang, C. H. *et al.* A retrospective study comparing tension-free vaginal tape and transobturator suburethral tape for surgical treatment of female stress urinary incontinence — a preliminary report. *J. Chin. Med. Assoc.* **70**, 541–544 (2007).

100. Davis, N. F., Smyth, L. G., Giri, S. K. & Flood, H. D. Evaluation of endoscopic laser excision of polypropylene mesh/sutures following anti-incontinence procedures. *J. Urol.* **188**, 1828–1832 (2012).
101. Deffieux, X. et al. Prepubic and thigh abscess after successive placement of two suburethral slings. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 571–574 (2007).
102. Elliott, J. E. & Maslow, K. D. Remote discovery of an asymptomatic bowel perforation by a mid-urethral sling. *Int. Urogynecol. J.* **23**, 243–245 (2012).
103. Feiner, B. & Maher, C. Vaginal mesh contraction: definition, clinical presentation, and management. *Obstet. Gynecol.* **115**, 325–330 (2010).
104. Fourie, T. & Cohen, P. L. Delayed bowel erosion by tension-free vaginal tape (TVT). *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **14**, 362–364 (2003).
105. Govier, F. et al. Complications of transvaginal silicone-coated polyester synthetic mesh sling. *Urology* **66**, 741–745 (2005).
106. Gruber, D. D., Wiersma, D. S., Dunn, J. S., Meldrum, K. A. & Krivak, T. C. Cecal perforation complicating placement of a transvaginal tension-free vaginal tape. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 671–673 (2007).
107. Hansen, B. L., Dunn, G. E., Norton, P., Hsu, Y. & Nygaard, I. Long-term follow-up of treatment for synthetic mesh complications. *Female Pelvic Med. Reconstr. Surg.* **20**, 126–130 (2014).
108. Hodroff, M., Portis, A. & Siegel, S. W. Endoscopic removal of intravaginal polypropylene sling with the holmium laser. *J. Urol.* **172**, 1361–1362 (2004).
109. Lee, J. K., Agnew, G. & Dwyer, P. L. Mesh-related chronic infections in silicone-coated polyester suburethral slings. *Int. Urogynecol. J.* **22**, 29–35 (2011).
110. Marcus-Braun, N., Bourret, A. & von Theobald, P. Persistent pelvic pain following transvaginal mesh surgery: a cause for mesh removal. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **162**, 224–228 (2012).
111. Marcus-Braun, N. & von Theobald, P. Mesh removal following transvaginal mesh placement: a case series of 104 operations. *Int. Urogynecol. J.* **21**, 423–430 (2010).
112. Vervest, H., Bongers, M. & van der Wurff, A. Nerve injury: an exceptional cause of pain after TVT. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **6**, 665–667 (2006).
113. Yeung, P. Jr, Sokol, A., Walton, B. & Iglesia, C. Thigh abscess mistaken for sarcoma following transobuturator tape: a case report and literature review. *J. Minim. Invasive Gynecol.* **14**, 657–659 (2007).
114. Hou, J. C., Alhalabi, F., Lemack, G. E. & Zimmer, P. E. Outcome of transvaginal mesh and tape removed for pain only. *J. Urol.* **192**, 856–860 (2014).
115. Bekelman, J. E., Li, Y. & Gross, C. P. Scope and impact of financial conflicts of interest in biomedical research: a systematic review. *JAMA* **289**, 454–465 (2003).
116. Petri, E. & Ashok, K. Comparison of late complications of retropubic and transobuturator slings in stress urinary incontinence. *Int. Urogynecol. J.* **23**, 321–325 (2012).
117. Barnes, N. M., Dmochowski, R. R., Park, R. & Nitti, V. W. Pubovaginal sling and pelvic prolapse repair in women with occult stress urinary incontinence: effect on postoperative emptying and voiding symptoms. *Urology* **59**, 856–860 (2002).
118. Sivanesan, K., Abdel-Fattah, M. & Tierney, J. Perineal cellulitis and persistent vaginal erosion after transobuturator tape (Obtape)—case report and review of the literature. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 219–221 (2007).
119. Errando, C., Batista, J. E. & Arano, P. Polytetrafluoroethylene sling for failure in female stress incontinence surgery. *World J. Urol.* **14**, S48–S50 (1996).
120. Connolly, T. P. Necrotizing surgical site infection after tension-free vaginal tape. *Obstet. Gynecol.* **104**, 1275–1276 (2004).
121. Beck, R. P. & Lai, A. R. Results in treating 88 cases of recurrent urinary stress incontinence with the Oxford fascia lata sling procedure. *Am. J. Obstet. Gynecol.* **142**, 649–651 (1982).
122. Neuman, M. Infected haematoma following tension-free vaginal tape implantation. *J. Urol.* **168**, 2549 (2002).
123. Agostini, A. et al. Abscess of the thigh and psoas muscle after transobuturator suburethral sling procedure. *Acta Obstet. Gynecol. Scand.* **85**, 628–629 (2006).
124. Domingo, S., Alama, P., Ruiz, N., Perales, A. & Pellicer, A. Diagnosis, management and prognosis of vaginal erosion after transobuturator suburethral tape procedure using a nonwoven thermally bonded polypropylene mesh. *J. Urol.* **173**, 1627–1630 (2005).
125. Siegel, A. L. Vaginal mesh extrusion associated with use of Mentor transobuturator sling. *Urology* **66**, 995–999 (2005).
126. Rafii, A., Jacob, D. & Deval, B. Obturator abscess after transobuturator tape for stress urinary incontinence. *Obstet. Gynecol.* **108**, 720–723 (2006).
127. Benassi, G., Marconi, L., Accorsi, F., Angeloni, M. & Benassi, L. Abscess formation at the ischiorectal fossa 7 months after the application of a synthetic transobuturator sling for stress urinary incontinence in a type II diabetic woman. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 697–699 (2007).
128. Babalola, E. O., Famuyide, A. O., McGuire, L. J., Gebhart, J. B. & Klingele, C. J. Vaginal erosion, sinus formation, and ischiorectal abscess following transobuturator tape: ObTape implantation. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **17**, 418–421 (2006).
129. Fialkow, M. F., Lentz, G. M., Miller, E. A. & Miller, J. L. Complications from transvaginal pubovaginal slings using bone anchor fixation. *Urology* **64**, 1127–1132 (2004).
130. Riedler, I. et al. Fournier's gangrene after tension-free vaginal tape (TVT) procedure. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **15**, 145–146 (2004).
131. Frederick, R. W., Carey, J. M. & Leach, G. E. Osseous complications after transvaginal bone anchor fixation in female pelvic reconstructive surgery: report from single largest prospective series and literature review. *Urology* **64**, 669–674 (2004).
132. Kascak, P. K. Fatal Injury of the small intestine during retropubic sling placement—a case report. *Obstet. Gynecol. Cases Rev.* **1** (2014).
133. Olagundoye, V. O., Shah, M. M. & Jackson, S. R. Delayed presentation of small bowel trauma during insertion of tension free vaginal tape (TVT) sling. *J. Obstet. Gynaecol.* **27**, 92–93 (2007).
134. Meschia, M., Busacca, M., Pifarotti, P. & De Marinis, S. Bowel perforation during insertion of tension-free vaginal tape (TVT). *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **13**, 263–265 (2002).
135. Gomelsky, A. & Dmochowski, R. R. Update on the management of overactive bladder: patient considerations and adherence. *Open Access J. Urol.* **3**, 7–17 (2010).
136. Wagg, A., Compion, G., Fahey, A. & Siddiqui, E. Persistence with prescribed antimuscarinic therapy for overactive bladder: a UK experience. *BJU Int.* **110**, 1767–1774 (2012).
137. Brostrom, S. & Hallas, J. Persistence of antimuscarinic drug use. *Eur. J. Clin. Pharmacol.* **65**, 309–314 (2009).
138. Roupert, M. et al. Laparoscopic surgical complete sling resection for tension-free vaginal tape-related complications refractory to first-line conservative management: a single-centre experience. *Eur. Urol.* **58**, 270–274 (2010).
139. Dindo, D., Demartines, N. & Clavien, P. Classification of surgical complications: a new proposal with evaluation in a cohort of 6,336 patients and results of a survey. *Ann. Surg.* **240**, 205–213 (2004).
140. Richter, H. E. et al. Retropubic versus transobuturator midurethral slings for stress incontinence. *N. Engl. J. Med.* **362**, 2066–2076 (2010).
141. The Urinary Incontinence Treatment Network. The trial of mid-urethral slings (TOMUS): design and methodology. *J. Appl. Res.* **8** (2008).
142. Strasberg, S. M., Linehan, D. C. & Hawkins, W. G. The accordancy severity grading system of surgical complications. *Ann. Surg.* **250**, 177–186 (2009).
143. Suskind, A. M. et al. Effectiveness of mesh compared with nonmesh sling surgery in Medicare beneficiaries. *Obstet. Gynecol.* **122**, 546–552 (2013).
144. Madsen, A. M. et al. A cohort study comparing a single-incision sling with a retropubic midurethral sling. *Int. Urogynecol. J.* **25**, 351–358 (2014).
145. Lord, H. E. et al. A randomized controlled equivalence trial of short-term complications and efficacy of tension-free vaginal tape and suprapubic urethral support sling for treating stress incontinence. *BJU Int.* **98**, 367–376 (2006).
146. Young, S. B., Howard, A. E. & Baker, S. P. Mersilene mesh sling: short- and long-term clinical and urodynamic outcomes. *Am. J. Obstet. Gynecol.* **185**, 32–40 (2001).
147. Hodroff, M. A., Sutherland, S. E., Kesha, J. B. & Siegel, S. W. Treatment of stress incontinence with the SPARC sling: intraoperative and early complications of 445 patients. *Urology* **66**, 760–762 (2005).
148. Palma, P. et al. Transobuturator SAFYRE sling is as effective as the transvaginal procedure. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **16**, 487–491 (2005).
149. Skala, C. et al. The IUGA/ICS classification of complications of prosthesis and graft insertion: a comparative experience in incontinence and prolapse surgery. *Int. Urogynecol. J.* **22**, 1429–1435 (2011).
150. Haylen, B. et al. An International Urogynecological Association (IUGA)/International Continence Society (ICS) joint terminology and classification of the complications related directly to the insertion of prosthesis (meshes, implants, tapes) and grafts in female pelvic floor surgery. *Neurourol. Urodynam.* **30**, 2–12 (2011).
151. Abouassaly, R. et al. Complications of tension-free vaginal tape surgery: a multi-institutional review. *BJU Int.* **94**, 110–113 (2004).
152. Jakimiuk, A. J. et al. Surgical treatment of stress urinary incontinence using the tension-free vaginal tape-obturator system (TVT-O) technique. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **135**, 127–131 (2007).

153. Kim, J. C. *et al.* A safety and quality of life analysis of intravaginal slingplasty in female stress incontinence: a prospective, open label, multicentre, and observational study. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 1331–1335 (2007).
154. Siddighi, S. & Karram, M. M. Surgical and nonsurgical approaches to treat voiding dysfunction following antiincontinence surgery. *Curr. Opin. Obstet. Gynecol.* **19**, 490–495 (2007).
155. Eandi, J. A., Tanaka, S. T., Henthall, N. J., O'Connor, R. C. & Stone, A. R. Self-reported urinary continence outcomes for repeat midurethral synthetic sling placement. *Int. Braz. J. Urol.* **34**, 336–342 (2008).
156. Tcherniakovsky, M., Fernandes, C. E., Bezerra, C. A., Del Roy, C. A. & Wroclawski, E. R. Comparative results of two techniques to treat stress urinary incontinence: synthetic transobturator and aponeurotic slings. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 961–966 (2009).
157. Paparella, R. *et al.* Prospective randomized trial comparing synthetic vs biological out-in transobturator tape: a mean 3-year follow-up study. *Int. Urogynecol. J.* **21**, 1327–1336 (2010).
158. Bandarian, M., Ghanbari, Z. & Asgari, A. Comparison of transobturator tape (TOT) vs Burch method in treatment of stress urinary incontinence. *J. Obstet. Gynaecol.* **31**, 518–520 (2011).
159. Chen, H. Y., Yeh, L. S., Chang, W. C. & Ho, M. Analysis of risk factors associated with surgical failure of inside-out transobturator vaginal tape for treating urodynamic stress incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 443–447 (2007).
160. Mellier, G., Mistrangelo, E., Gery, L., Philippe, C. & Patrice, M. Tension-free obturator tape (Monarc Subfascial Hammock) in patients with or without associated procedures. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 165–172 (2007).
161. Tinelli, A. *et al.* Presurgical promestriene therapy in postmenopausal women with stress urinary incontinence. *Gynecol. Endocrinol.* **23**, 445–450 (2007).
162. Withagen, M. I. & Milani, A. L. Which factors influenced the result of a tension free vaginal tape operation in a single teaching hospital? *Acta Obstet. Gynecol. Scand.* **86**, 1136–1139 (2007).
163. Aniulienė, R. Tension-free vaginal tape versus tension-free vaginal tape obturator (inside-outside) in the surgical treatment of female stress urinary incontinence. *Medicina (Kaunas)* **45**, 639–643 (2009).
164. Su, T. H. *et al.* Tension-free vaginal tape-obturator procedure for treatment of severe urodynamic stress incontinence: subjective and objective outcomes during 2 years of follow-up. *J. Obstet. Gynaecol. Res.* **35**, 1077–1082 (2009).
165. Lau, H. H., Su, T. H., Su, C. H., Lee, M. Y. & Sun, F. J. Short-term impact of tension-free vaginal tape obturator procedure on sexual function in women with stress urinary incontinence. *J. Sex. Med.* **7**, 1578–1584 (2010).
166. Palva, K. *et al.* A randomized trial comparing tension-free vaginal tape with tension-free vaginal tape-obturator: 36-month results. *Int. Urogynecol. J.* **21**, 1049–1055 (2010).
167. Lleberia-Juanos, J. *et al.* De novo urgency after tension-free vaginal tape versus transobturator tape procedure for stress urinary incontinence. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **155**, 229–232 (2011).
168. Freeman, R. *et al.* What patients think: patient-reported outcomes of retropubic versus trans-obturator mid-urethral slings for urodynamic stress incontinence—a multi-centre randomised controlled trial. *Int. Urogynecol. J.* **22**, 279–286 (2011).
169. Liu, P. E. *et al.* Outcome of tension-free obturator tape procedures in obese and overweight women. *Int. Urogynecol. J.* **22**, 259–263 (2011).
170. Jeon, M. J. *et al.* Surgical therapeutic index of tension-free vaginal tape and transobturator tape for stress urinary incontinence. *Gynecol. Obstet. Invest.* **65**, 41–46 (2008).
171. Castillo-Pino, E., Sasson, A. & Pons, J. E. Comparison of retropubic and transobturator tension-free vaginal implants for the treatment of stress urinary incontinence. *Int. J. Gynaecol. Obstet.* **110**, 23–26 (2010).
172. Cornu, J. N. *et al.* Midterm prospective evaluation of TVT-Secur reveals high failure rate. *Eur. Urol.* **58**, 157–161 (2010).
173. Lo, T. S. Combined pelvic reconstructive surgery and transobturator tape (monarc) in women with advanced prolapse and urodynamic stress incontinence: a case control series. *J. Minim. Invasive Gynecol.* **16**, 163–168 (2009).
174. Bianchi-Ferraro, A. M. *et al.* Single-incision sling compared with transobturator sling for treating stress urinary incontinence: a randomized controlled trial. *Int. Urogynecol. J.* **24**, 1459–1465 (2013).
175. Wadie, B. S., Mansour, A., El-Hefnawy, A. S., Nabeeh, A. & Khair, A. A. Minimum 2-year follow-up of mid-urethral slings, effect on quality of life, incontinence impact and sexual function. *Int. Urogynecol. J.* **21**, 1485–1490 (2010).
176. Wu, J. Y., He, H. C., Chen, S. W., Jin, X. D. & Zhou, Y. X. Surgical therapies of female stress urinary incontinence: experience in 228 cases. *Int. Urogynecol. J.* **21**, 645–649 (2010).
177. Sabadell, J. *et al.* Usefulness of retropubic tape for recurrent stress incontinence after transobturator tape failure. *Int. Urogynecol. J.* **22**, 1543–1547 (2011).
178. Masata, J. *et al.* Randomized trial of a comparison of the efficacy of TVT-O and single-incision tape TVT SECUR systems in the treatment of stress urinary incontinent women—2-year follow-up. *Int. Urogynecol. J.* **23**, 1403–1412 (2012).
179. Dyrkorn, O. A., Kulseng-Hanssen, S. & Sandvik, L. TVT compared with TVT-O and TOT: results from the Norwegian National Incontinence Registry. *Int. Urogynecol. J.* **21**, 1321–1326 (2010).
180. Liapis, A., Bakas, P. & Creatsas, G. Efficacy of inside-out transobturator vaginal tape (TVTO) at 4 years follow up. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **148**, 199–201 (2010).
181. Andrada Hamer, M., Larsson, P. G., Teleman, P., Bergqvist, C. E. & Persson, J. One-year results of a prospective randomized, evaluator-blinded, multicentre study comparing TVT and TVT Secur. *Int. Urogynecol. J.* **24**, 223–229 (2013).
182. Poza, J. L. *et al.* Trans-obturator suburethral tape for female stress incontinence: a cohort of 254 women with 1-year to 2-year follow-up. *Acta Obstet. Gynecol. Scand.* **87**, 232–239 (2008).
183. Rechberger, T., Futyma, K., Jankiewicz, K., Adamiak, A. & Skorpupski, P. The clinical effectiveness of retropubic (IVS-02) and transobturator (IVS-04) midurethral slings: randomized trial. *Eur. Urol.* **56**, 24–30 (2009).
184. Sun, M. J. & Tsai, H. D. Is transobturator suburethral sling effective for treating female urodynamic stress incontinence with low maximal urethral closure pressure? *Taiwan. J. Obstet. Gynecol.* **50**, 20–24 (2011).
185. Sakalis, V. I., Gkotsi, A. C., Triantafyllidis, A., Giouris, A. & Charalambous, S. Transurethral holmium laser intravesical tape excision following TVT procedure: results from seven patients in a 12-month follow-up. *Int. Urogynecol. J.* **23**, 769–777 (2012).
186. Palva, K. & Nilsson, C. G. Prevalence of urinary urgency symptoms decreases by mid-urethral sling procedures for treatment of stress incontinence. *Int. Urogynecol. J.* **22**, 1241–1247 (2011).
187. Marques, A. L., Aparicio, C. & Negrao, L. Perineal-cellulitis as a late complication of trans-obturator sub-urethral tape, Obtape. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 821–822 (2007).
188. Goldman, H. B. Large thigh abscess after placement of synthetic transobturator sling. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **17**, 295–296 (2006).
189. Mahajan, S. T., Kenton, K., Bova, D. A. & Brubaker, L. Transobturator tape erosion associated with leg pain. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **17**, 66–68 (2006).
190. Amid, P. K. Classification of biomaterials and their related complications in abdominal wall hernia surgery. *Hernia* **1**, 15–21 (1997).
191. Weinberger, J. M., Purohit, R. S. & Blaivas, J. G. Mesh infection of a male sling. *J. Urol.* **190**, 1054–1055 (2013).
192. Johnson, D. W., ElHajj, M., O'Brien-Best, E. L., Miller, H. J. & Fine, P. M. Necrotizing fasciitis after tension-free vaginal tape (TVT) placement. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **14**, 291–293 (2003).
193. Busby, G. & Broome, J. Necrotising fasciitis following unrecognized bladder injury during transobturator sling procedure. *BJOG* **114**, 111–112 (2007).
194. Wadie, B. S., Edwan, A. & Nabeeh, A. M. Autologous fascial sling vs polypropylene tape at short-term followup: a prospective randomized study. *J. Urol.* **174**, 990–993 (2005).
195. Kuo, H. C. Long-term surgical results of pubovaginal sling procedure using polypropylene mesh in the treatment of stress urinary incontinence. *Urol. Int.* **74**, 147–152 (2005).
196. Kuuva, N. & Nilsson, C. G. A nationwide analysis of complications associated with the tension-free vaginal tape (TVT) procedure. *Acta Obstet. Gynecol. Scand.* **81**, 72–77 (2002).
197. Siegel, A. L., Kim, M., Goldstein, M., Levey, S. & Ilbeigi, P. High incidence of vaginal mesh extrusion using the intravaginal slingplasty sling. *J. Urol.* **174**, 1308–1311 (2005).
198. Choe, J. M., Ogan, K. & Bennett, S. Antibacterial mesh sling: a prospective outcome analysis. *Urology* **55**, 515–520 (2000).
199. Hom, D., Desautel, M. G., Lumerman, J. H., Feraren, R. E. & Badlani, G. H. Pubovaginal sling using polypropylene mesh and Vesica bone anchors. *Urology* **51**, 708–713 (1998).
200. Tseng, L. H., Wang, A. C., Lin, Y. H., Li, S. J. & Ko, Y. J. Randomized comparison of the suprapubic arc sling procedure vs tension-free vaginal taping for stress incontinent women. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **16**, 230–235 (2005).
201. Ashok, K. & Petri, E. Failures and complications in pelvic floor surgery. *World J. Urol.* **30**, 487–494 (2012).
202. Onol, F. F. *et al.* Minimum 1.5-year results of “surgeon-tailored” transvaginal mesh repair for female stress urinary incontinence and pelvic organ prolapse. *Urology* **80**, 273–279 (2012).
203. Naidu, A. *et al.* Transobturator tape for stress incontinence: the North Queensland experience. *Aust. N. Z. J. Obstet. Gynaecol.* **45**, 446–449 (2005).

204. Youn, C. S., Shin, J. H. & Na, Y. G. Comparison of TOA and TOT for treating female stress urinary incontinence: short-term outcomes. *Korean J. Urol.* **51**, 544–549 (2010).
205. Takahashi, S. *et al.* Transvaginal mesh (TVM) reconstruction with TVT/TOT sling for vaginal prolapse concurrent with stress urinary incontinence. *Aktuelle Urol.* **41**, S20–S23 (2010).
206. Schimpf, M. O. *et al.* Sling surgery for stress urinary incontinence in women: a systematic review and metaanalysis. *Am. J. Obstet. Gynecol.* **211**, 71.e1–e71 (2014).
207. Brito, L. M. *et al.* Comparison of the outcomes of the sling technique using a commercial and hand-made polypropylene sling. *Int. Braz. J. Urol.* **37**, 519–527 (2011).
208. Chen, X., Li, H., Fan, B., Yang, X. & Tong, X. An inexpensive modified transobturator vaginal tape inside-out procedure for the surgical treatment of female stress urinary incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 1365–1368 (2009).
209. Chene, G. *et al.* Long-term results of tension-free vaginal tape (TVT) for the treatment of female urinary stress incontinence. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **134**, 87–94 (2007).
210. Altman, D., Anzen, B., Morlin, B. & Falconer, C. Tension-free vaginal tape after failed cystoscopic transurethral injections for stress urinary incontinence. *Scand. J. Urol. Nephrol.* **43**, 506–508 (2009).
211. Domingo, S. *et al.* Transobturator tape procedure outcome: a clinical and quality of life analysis of a 1-year follow-up. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 895–900 (2007).
212. Glavind, K. & Larsen, T. Long-term follow-up of intravaginal slingplasty operation for urinary stress incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 1081–1083 (2008).
213. Gold, R. S., Grouz, A., Pauzner, D., Lessing, J. & Gordon, D. Bladder perforation during tension-free vaginal tape surgery: does it matter? *J. Reprod. Med.* **52**, 616–618 (2007).
214. Guerrero, K. L. *et al.* A randomised controlled trial comparing TVT, Pelvicol and autologous fascial slings for the treatment of stress urinary incontinence in women. *BJOG* **117**, 1493–1502 (2010).
215. Haliloglu, B., Karateke, A., Coksuer, H., Peker, H. & Cam, C. The role of urethral hypermobility and intrinsic sphincteric deficiency on the outcome of transobturator tape procedure: a prospective study with 2-year follow-up. *Int. Urogynecol. J.* **21**, 173–178 (2010).
216. Hellberg, D., Holmgren, C., Lanner, L. & Nilsson, S. The very obese woman and the very old woman: tension-free vaginal tape for the treatment of stress urinary incontinence. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 423–429 (2007).
217. Jijon, A., Hegde, A., Arias, B., Aguilar, V. & Davila, G. W. An inelastic retropubic suburethral sling in women with intrinsic sphincter deficiency. *Int. Urogynecol. J.* **24**, 1325–1330 (2013).
218. Juma, S. & Brito, C. G. Transobturator tape (TOT): Two years follow-up. *Neurourol. Urodyn.* **26**, 37–41 (2007).
219. Kang, M. B., Kim, H. G., Paick, S. H., Lho, Y. S. & Park, H. K. Intraoperative maximal urethral closing pressure measurement: a new technique of tape tension adjustment in transobturator sling surgery? *Int. Braz. J. Urol.* **37**, 751–757 (2011).
220. Karateke, A., Haliloglu, B., Cam, C. & Sakalli, M. Comparison of TVT and TVT-O in patients with stress urinary incontinence: short-term cure rates and factors influencing the outcome. A prospective randomised study. *Aust. N. Z. J. Obstet. Gynaecol.* **49**, 99–105 (2009).
221. Kim, J., Lucioni, A., Govier, F. & Kobashi, K. Worse long-term surgical outcomes in elderly patients undergoing SPARC retropubic midurethral sling placement. *BJU Int.* **108**, 708–712 (2011).
222. Lee, K. S. *et al.* Outcomes following repeat mid urethral synthetic sling after failure of the initial sling procedure: rediscovery of the tension-free vaginal tape procedure. *J. Urol.* **178**, 1370–1374 (2007).
223. Lier, D. *et al.* Trans-obturator tape compared with tension-free vaginal tape in the surgical treatment of stress urinary incontinence: a cost utility analysis. *BJOG* **118**, 550–556 (2011).
224. Lim, Y. N. *et al.* Do the Advantage slings work as well as the tension-free vaginal tapes? *Int. Urogynecol. J.* **21**, 1157–1162 (2010).
225. Pace, G. & Vicentini, C. Female sexual function evaluation of the tension-free vaginal tape (TVT) and transobturator suburethral tape (TOT) incontinence surgery: results of a prospective study. *J. Sex. Med.* **5**, 387–393 (2008).
226. Paick, J. S., Oh, S. J., Kim, S. W. & Ku, J. H. Tension-free vaginal tape, suprapubic arc sling, and transobturator tape in the treatment of mixed urinary incontinence in women. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 123–129 (2008).
227. Reich, A., Kohorst, F., Kreienberg, R. & Flock, F. Long-term results of the tension-free vaginal tape procedure in an unselected group: a 7-year follow-up study. *Urology* **78**, 774–777 (2011).
228. Shaker, H. S., Ban, H. M., Hegazy, A. S. & Mansour, M. F. Functional and quality of life outcome of transobturator tape for treatment of female stress urinary. *Int. Urogynecol. J.* **22**, 99–103 (2011).
229. Tommaselli, G. A. *et al.* Efficacy and safety of TVT-O and TVT-Secur in the treatment of female stress urinary incontinence: 1-year follow-up. *Int. Urogynecol. J.* **21**, 1211–1217 (2010).
230. Trabuco, E. C. *et al.* Medium-term comparison of continence rates after rectus fascia or midurethral sling placement. *Am. J. Obstet. Gynecol.* **200**, 300.e1–e6 (2009).
231. Wadie, B. S. & El-Hefnawy, A. S. TVT versus TOT, 2-year prospective randomized study. *World J. Urol.* **31**, 645–649 (2013).
232. Wang, Y. J. *et al.* Comparison of three mid-urethral tension-free tapes (TVT, TVT-O, and TVT-Secur) in the treatment of female stress urinary incontinence: 1-year follow-up. *Int. Urogynecol. J.* **22**, 1369–1374 (2011).
233. Yang, X. *et al.* TVT-O vs. TVT for the treatment of SUL: a non-inferiority study. *Int. Urogynecol. J.* **23**, 99–104 (2012).
234. Yoon, C. J. & Jung, H. C. Three-year outcomes of the innovative replacement of incontinence surgery procedure for treatment of female stress urinary incontinence: comparison with tension-free vaginal tape procedure. *J. Korean Med. Sci.* **22**, 497–501 (2007).
235. Zuger, V. *et al.* TVT vs. TOT: a comparison in terms of continence results, complications and quality of life after a median follow-up of 48 months. *Int. Urol. Nephrol.* **42**, 915–920 (2010).
236. Huffaker, R. K., Yandell, P. M. & Shull, B. L. Tension-free vaginal tape bowel perforation. *Int. Urogynecol. J.* **21**, 251–253 (2010).
237. Phillips, L., Flood, C. G. & Schulz, J. A. Case report of tension-free vaginal tape-associated bowel obstruction and relationship to body habitus. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 367–368 (2009).
238. Agostini, A. *et al.* Immediate complications of tension-free vaginal tape (TVT): results of a French survey. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **124**, 237–239 (2006).
239. Arrabal-Polo, M. *et al.* Complications from the placement of a tension-free suburethral sling using the transobturator and retropubic methods for treatment of female urinary incontinence. *Urol. Int.* **89**, 97–102 (2012).
240. Minaglia, S., Ozel, B., Klutke, C., Ballard, C. & Klutke, J. Bladder injury during transobturator sling. *Urology* **64**, 376–377 (2004).
241. Klutke, C. *et al.* Urinary retention after tension-free vaginal tape procedure: incidence and treatment. *Urology* **58**, 697–701 (2001).
242. Niemczyk, P., Klutke, J. J., Carlin, B. I. & Klutke, C. G. United States experience with tension-free vaginal tape procedure for urinary stress incontinence: assessment of safety and tolerability. *Tech. Urol.* **7**, 261–265 (2001).
243. Blaivas, J. G. & Purohit, R. S. Post-traumatic female urethral reconstruction. *Curr. Urol. Rep.* **9**, 397–404 (2008).
244. Goldman, H. B. Simple sling incision for the treatment of iatrogenic urethral obstruction. *Urology* **62**, 714–718 (2003).
245. Angioli, R. *et al.* Suburethral sling in autoimmune patients: complications, quality of life, and success rate. *Int. Urogynecol. J.* **23**, 453–457 (2012).
246. Barber, M. D. *et al.* Risk factors associated with failure 1 year after retropubic or transobturator midurethral slings. *Am. J. Obstet. Gynecol.* **199**, 666.e1–e7 (2008).
247. Djehdian, L. M. *et al.* Transobturator sling compared with single-incision mini-sling for the treatment of stress urinary incontinence: a randomized controlled trial. *Obstet. Gynecol.* **123**, 553–561 (2014).
248. Goktolga, U. *et al.* Tension-free vaginal tape for surgical relief of intrinsic sphincter deficiency: results of 5-year follow-up. *J. Minim. Invasive Gynecol.* **15**, 78–81 (2008).
249. Kociszewski, J. *et al.* Tape functionality: position, change in shape, and outcome after TVT procedure—mid-term results. *Int. Urogynecol. J.* **21**, 795–800 (2010).
250. Latthe, P. M., Patodi, M. & Constantine, G. Transobturator tape procedure in stress urinary incontinence: UK experience of a district general hospital. *J. Obstet. Gynaecol.* **27**, 177–180 (2007).
251. Sander, P., Sorensen, F. & Lose, G. Does the tension-free vaginal tape procedure (TVT) affect the voiding function over time? Pressure-flow studies 1 year and 3(1/2) years after TVT. *Neurourol. Urodyn.* **26**, 995–997 (2007).
252. Sivaslioglu, A. A., Unlubilgin, E., Aydogmus, S., Keskin, L. & Dolen, I. A prospective randomized controlled trial of the transobturator tape and tissue fixation mini-sling in patients with stress urinary incontinence: 5-year results. *J. Urol.* **188**, 194–199 (2012).
253. Smith, T. *et al.* Pathologic evaluation of explanted vaginal mesh: interdisciplinary experience from a referral centre. *Female Pelvic Med. Reconstr. Surg.* **19**, 238–241 (2013).
254. Tanuri, A. L. *et al.* [Retropubic and transobturator sling in treatment of stress urinary incontinence]. *Rev. Assoc. Med. Bras.* **56**, 348–354 (2010).
255. Tommaselli, G. A. *et al.* Tension-free vaginal tape-O and -Secur for the treatment of stress urinary incontinence: a thirty-six-month follow-up single-blind, double-arm, randomized study. *J. Minim. Invasive Gynecol.* **20**, 198–204 (2013).

256. Giberti, C., Gallo, F., Cortese, P. & Schenone, M. The suburethral tension adjustable sling (REMEEX system) in the treatment of female urinary incontinence due to 'true' intrinsic sphincter deficiency: results after 5 years of mean follow-up. *BJU Int.* **108**, 1140–1144 (2011).
257. Heinonen, P., Ala-Nissila, S., Raty, R., Laurikainen, E. & Kiilholma, P. Objective cure rates and patient satisfaction after the transobturator tape procedure during 6.5-year follow-up. *J. Minim. Invasive Gynecol.* **20**, 73–78 (2013).
258. Kim, J. H., Doo, S. W., Yang, W. J. & Song, Y. S. Laparoscopic transvesical excision and reconstruction in the management of mid-urethral tape mesh erosion and stones around the bladder neck: initial experiences. *BJU Int.* **110**, E1009–E1013 (2012).
259. Sivaslioglu, A. A., Unlubgilin, E. & Dolen, I. The multifilament polypropylene tape erosion trouble: tape structure vs surgical technique. Which one is the cause? *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 417–420 (2008).
260. Ward, K. L., Hilton, P. & The UK and Ireland TVT trial group. Tension-free vaginal tape versus colposuspension for primary urodynamic stress incontinence: 5-year follow up. *BJOG* **115**, 226–233 (2008).
261. Tjeldink, M. M., Vierhout, M. E., Heesakkers, J. P. & Withagen, M. I. Surgical management of mesh-related complications after prior pelvic floor reconstructive surgery with mesh. *Int. Urogynecol. J.* **22**, 1395–1404 (2011).
262. Abdel-fattah, M., Mostafa, A., Young, D. & Ramsay, I. Evaluation of transobturator tension-free vaginal tapes in the management of women with mixed urinary incontinence: one-year outcomes. *Am. J. Obstet. Gynecol.* **205**, 150.e1–e6 (2011).
263. Ballester, M. et al. Four-year functional results of the suburethral sling procedure for stress urinary incontinence: a French prospective randomized multicentre study comparing the retropubic and transobturator routes. *World J. Urol.* **30**, 117–122 (2012).
264. Botros, S. M. et al. Detrusor overactivity and urge urinary incontinence following trans obturator versus midurethral slings. *Neurourol. Urodyn.* **26**, 42–45 (2007).
265. But, I. & Pakiz, M. Irritative symptoms are the main predictor of satisfaction rate in women after transobturator tape procedures. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 791–796 (2009).
266. Groutz, A. et al. The safety and efficacy of the "inside-out" trans-obturator TVT in elderly versus younger stress-incontinent women: A prospective study of 353 consecutive patients. *Neurourol. Urodyn.* **30**, 380–383 (2011).
267. Lee, H. N. et al. Transurethral injection of bulking agent for treatment of failed mid-urethral sling procedures. *Int. Urogynecol. J.* **21**, 1479–1483 (2010).
268. Maroto, J. R. et al. Transobturator adjustable tape (TOA) permits to correct postoperatively the tension applied in stress incontinence surgery. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 797–805 (2009).
269. Holmgren, C., Nilsson, S., Lanner, L. & Hellberg, D. Frequency of de novo urgency in 463 women who had undergone the tension-free vaginal tape (TVT) procedure for genuine stress urinary incontinence—a long-term follow-up. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **132**, 121–125 (2007).
270. Stewart, W. F. et al. Prevalence and burden of overactive bladder in the United States. *World J. Urol.* **20**, 327–336 (2003).
271. Sweeney, D. D. & Leng, W. W. Treatment of postoperative voiding dysfunction following incontinence surgery. *Curr. Urol. Rep.* **6**, 365–370 (2005).
272. Dmochowski, R., Scarpero, H. M. & Starkman, J. S. in *Campbell-Walsh Urology* (eds Wein, A. J., Kavoussi, L. R., Novick, A. C., Partin, A. W. & Peters, C. A.) 2251–2272 (Saunders, Philadelphia, 2007).
273. Aponte, M. M. et al. Urodynamics for clinically suspected obstruction after anti-incontinence surgery in women. *J. Urol.* **190**, 598–602 (2013).
274. Ozel, B., Minaglia, S., Hurtado, E., Klutke, C. G. & Klutke, J. J. Treatment of voiding dysfunction after transobturator tape procedure. *Urology* **64**, 1030 (2004).
275. Kawasaki, A. et al. Comparing the risk of urethrolisis for the treatment of voiding dysfunction between two retropubic mesh slings: a case-control study. *Int. Urogynecol. J.* **24**, 589–594 (2013).
276. Ridgeway, B. et al. Early experience with mesh excision for adverse outcomes after transvaginal mesh placement using prolapse kits. *Am. J. Obstet. Gynecol.* **199**, 703.e1–e7 (2008).
277. Reynolds, W. S. et al. Obturator foramen dissection for excision of symptomatic transobturator mesh. *J. Urol.* **187**, 1680–1684 (2012).
278. Cetinel, B. & Tarcan, T. Management of complications after tension-free midurethral slings. *Korean J. Urol.* **54**, 651–659 (2013).
279. Roumeguere, T. et al. Trans-obturator vaginal tape (TOT) for female stress incontinence: one year follow-up in 120 patients. *Eur. Urol.* **48**, 805–809 (2005).
280. Long, C. Y. et al. Lateral excision of tension-free vaginal tape for the treatment of iatrogenic urethral obstruction. *Obstet. Gynecol.* **104**, 1270–1274 (2004).
281. Karra, M. M., Segal, J. L., Vassallo, B. J. & Kleeman, S. D. Complications and untoward effects of the tension-free vaginal tape procedure. *Obstet. Gynecol.* **101**, 929–932 (2003).
282. Petrou, S. P., Brown, J. A. & Blaivas, J. G. Suprameatal transvaginal urethrolisis. *J. Urol.* **161**, 1268–1271 (1999).
283. Carr, L. K. & Webster, G. D. Voiding dysfunction following incontinence surgery: diagnosis and treatment with retropubic or vaginal urethrolisis. *J. Urol.* **157**, 821–823 (1997).
284. Abed, H. et al. Incidence and management of graft erosion, wound granulation, and dyspareunia following vaginal prolapse repair with graft materials: a systematic review. *Int. Urogynecol. J.* **22**, 789–798 (2011).
285. Hinoul, P., Bonnet, P., Krofta, L., Waltregny, D. & de Leval, J. An anatomic comparison of the original versus a modified inside-out transobturator procedure. *Int. Urogynecol. J.* **22**, 997–1004 (2011).
286. Chen, X. et al. A modified inexpensive transobturator vaginal tape inside-out procedure versus tension-free vaginal tape for the treatment of SUI: a prospective comparative study. *Arch. Gynecol. Obstet.* **284**, 1461–1466 (2011).
287. Kristensen, I. et al. Complications of the tension-free vaginal tape procedure for stress urinary incontinence. *Int. Urogynecol. J.* **21**, 1353–1357 (2010).
288. Canete, P., Ortiz, E., Domingo, S. & Cano, A. Transobturator suburethral tape in the treatment of stress urinary incontinence: efficacy and quality of life after 5 year follow up. *Maturitas* **74**, 166–171 (2013).
289. Darai, E. et al. Functional results after the suburethral sling procedure for urinary stress incontinence: a prospective randomized multicentre study comparing the retropubic and transobturator routes. *Eur. Urol.* **51**, 795–801 (2007).
290. Hota, L. S. et al. TVT-Secur (Hammock) versus TVT-Obturator: a randomized trial of suburethral sling operative procedures. *Female Pelvic Med. Reconstr. Surg.* **18**, 41–45 (2012).
291. Mesens, T., Aich, A. & Bhal, P. S. Late erosions of mid-urethral tapes for stress urinary incontinence—need for long-term follow-up? *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 1113–1114 (2007).
292. Rapp, D. E., Govier, F. E. & Kobashi, K. C. Outcomes following mid-urethral sling placement in patients with intrinsic sphincteric deficiency: comparison of Sparc and Monarc slings. *Int. Braz. J. Urol.* **35**, 68–75 (2009).
293. Tommaselli, G. A. et al. Efficacy of a modified technique for TVT-O positioning: a twelve-month, randomized, single-blind, multicentre, non-inferiority study. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **167**, 225–229 (2013).
294. Klosterhalfen, B. & Klinge, U. Retrieval study at 623 human mesh explants made of polypropylene—impact of mesh class and indication for mesh removal on tissue reaction. *J. Biomed. Mater. Res. B. Appl. Biomater.* **101**, 1393–1399 (2013).
295. Yazdankhah Kenary, A. et al. Randomized clinical trial comparing lightweight mesh with heavyweight mesh for primary inguinal hernia repair. *Hernia* **17**, 471–477 (2013).
296. Khan, L. R., Kumar, S. & Nixon, S. J. Early results for new lightweight mesh in laparoscopic totally extra-peritoneal inguinal hernia repair. *Hernia* **10**, 303–308 (2006).
297. Peeters, E. et al. Sperm motility after laparoscopic inguinal hernia repair with lightweight meshes: 3-year follow-up of a randomised clinical trial. *Hernia* **18**, 361–367 (2014).
298. Nikkolo, C. et al. Three-year results of randomised clinical trial comparing lightweight mesh with heavyweight mesh for inguinal hernioplasty. *Hernia* **16**, 555–559 (2012).
299. Hirose, T. et al. Randomized clinical trial comparing lightweight or heavyweight mesh for mesh plug repair of primary inguinal hernia. *Hernia* **18**, 213–219 (2014).
300. Uzzaman, M. M., Ratnasingham, K. & Ashraf, N. Meta-analysis of randomized controlled trials comparing lightweight and heavyweight mesh for Lichtenstein inguinal hernia repair. *Hernia* **16**, 505–518 (2012).
301. Yamada, B., Govier, F., Stefanovic, K. & Kobashi, K. High rate of vaginal erosions associated with the mentor ObTape. *J. Urol.* **176**, 651–654 (2006).
302. Al-Badr, A. & Fouda, K. Suprapubic-assisted cystoscopic excision of intravesical tension-free vaginal tape. *J. Minim. Invasive Gynecol.* **12**, 370–371 (2005).
303. Cornel, E. B. & Vervest, H. A. Removal of a missed polypropylene tape by a combined transurethral and transabdominal endoscopic approach. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **16**, 247–249 (2005).
304. Baracat, F., Mitre, A. I., Kanashiro, H. & Montellato, N. I. Endoscopic treatment of vesical and urethral perforations after tension-free vaginal tape (TVT) procedure for female stress urinary incontinence. *Clinics (Sao Paulo)* **60**, 397–400 (2005).
305. Rosenblatt, P., Pulliam, S., Edwards, R. & Boyles, S. H. Suprapubically assisted operative cystoscopy in the management of intravesical TVT synthetic mesh segments. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **16**, 509–511 (2005).

306. Parekh, M. H., Minassian, V. A. & Poplawsky, D. Bilateral bladder erosion of a transobturator tape mesh. *Obstet. Gynecol.* **108**, 713–715 (2006).
307. Quiroz, L. H. & Cundiff, G. W. Transurethral resection of tension-free vaginal tape under tactile traction. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 873–875 (2009).
308. Wijffels, S. A., Elzevier, H. W. & Lycklama a Nijeholt, A. A. Transurethral mesh resection after urethral erosion of tension-free vaginal tape: report of three cases and review of literature. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 261–263 (2009).
309. Huwyler, M., Springer, J., Kessler, T. M. & Burkhard, F. C. A safe and simple solution for intravesical tension-free vaginal tape erosion: removal by standard transurethral resection. *BJU Int.* **102**, 582–585 (2008).
310. Oh, T. H. & Ryu, D. S. Transurethral resection of intravesical mesh after midurethral sling procedures. *J. Endourol.* **23**, 1333–1337 (2009).
311. Mendonca, T. M., Martinho, D. & Dos Reis, J. P. Late urethral erosion of transobturator suburethral mesh (Obtape): a minimally invasive management under local anaesthesia. *Int. Urogynecol. J.* **22**, 37–39 (2011).
312. Rogo-Gupta, L., Huynh, L., Hartshorn, T. G., Rodriguez, L. V. & Raz, S. Long-term symptom improvement and overall satisfaction after prolapse and incontinence graft removal. *Female Pelvic Med. Reconstr. Surg.* **19**, 352–355 (2013).
313. Ordorica, R., Rodriguez, A. R., Coste-Delvecchio, F., Hoffman, M. & Lockhart, J. Disabling complications with slings for managing female stress urinary incontinence. *BJU Int.* **102**, 333–336 (2008).
314. Blaivas, J. G. et al. Management of urethral stricture in women. *J. Urol.* **188**, 1778–1782 (2012).
315. Garcia-Urena, M. A. et al. Differences in polypropylene shrinkage depending on mesh position in an experimental study. *Am. J. Surg.* **193**, 538–542 (2007).
316. Klinge, U., Klosterhalfen, B., Müller, M., Ottinger, A. & Schumpelick, V. Shrinking of polypropylene mesh in vivo: an experimental study in dogs. *Eur. J. Surg.* **164**, 965–969 (1998).
317. Silva-Filho, A. L., Candido, E. B., Noronha, A. & Triginelli, S. A. Comparative study of autologous pubovaginal sling and synthetic transobturator (TOT) SAFYRE sling in the treatment of stress urinary incontinence. *Arch. Gynecol. Obstet.* **273**, 288–292 (2006).
318. Mamy, L. et al. Correlation between shrinkage and infection of implanted synthetic meshes using an animal model of mesh infection. *Int. Urogynecol. J.* **22**, 47–52 (2011).
319. Jacquetin, B. & Cosson, M. Complications of vaginal mesh: our experience. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 893–896 (2009).
320. Al-Wadi, K. & Al-Badr, A. Martius graft for the management of tension-free vaginal tape vaginal erosion. *Obstet. Gynecol.* **114**, 489–491 (2009).
321. Kokanali, M. et al. Risk factors for mesh erosion after vaginal sling procedures for urinary incontinence. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **177**, 146–150 (2014).
322. Mohr, S., Kuhn, P., Mueller, M. D. & Kuhn, A. Painful love-“hispareunia” after sling erosion of the female partner. *J. Sex. Med.* **8**, 1740–1746 (2011).
323. Kobashi, K. C. & Govier, F. E. Management of vaginal erosion of polypropylene mesh slings. *J. Urol.* **169**, 2242–2243 (2003).
324. Zoorob, D. & Karram, M. Management of mesh complications and vaginal constriction: a urogynecology perspective. *Urol. Clin. North Am.* **39**, 413–418 (2012).
325. Marks, B. K. & Goldman, H. B. Controversies in the management of mesh-based complications: a urology perspective. *Urol. Clin. North Am.* **39**, 419–428 (2012).
326. Lee, D., Dillon, B., Lemack, G., Gomelsky, A. & Zimmern, P. Transvaginal mesh kits—how “serious” are the complications and are they reversible? *Urology* **81**, 43–48 (2013).
327. Nicolson, A. & Adeyemo, D. Colovaginal fistula: a rare long-term complication of polypropylene mesh sacrocolpopexy. *J. Obstet. Gynaecol.* **29**, 444–445 (2009).
328. Hopkins, M. P. & Rooney, C. Entero mesh vaginal fistula secondary to abdominal sacral colpopexy. *Obstet. Gynecol.* **103**, 1035–1036 (2004).
329. Baumgartner, U., Magerl, W., Klein, T., Hopf, H. C. & Treede, R. D. Neurogenic hyperalgesia versus painful hypoalgesia: two distinct mechanisms of neuropathic pain. *Pain* **96**, 141–151 (2002).
330. Parnell, B. A., Johnson, E. A. & Zolnoun, D. A. Genitofemoral and perineal neuralgia after transobturator midurethral sling. *Obstet. Gynecol.* **119**, 428–431 (2012).
331. Anger, J. T. et al. Short-term outcomes of vaginal mesh placement among female Medicare beneficiaries. *Urology* **83**, 768–773 (2014).
332. Olah, K. S. Adductor compartment syndrome: An unusual complication of the trans-obturator tape procedure. *J. Obstet. Gynaecol.* **28**, 363–364 (2008).
333. Roth, T. M. Management of persistent groin pain after transobturator slings. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **18**, 1371–1373 (2007).
334. Daneshgari, F., Kong, W. & Swartz, M. Complications of mid urethral slings: important outcomes for future clinical trials. *J. Urol.* **180**, 1890–1897 (2008).
335. Sahin, A. F., Ilbey, Y. O. & Sahin, N. Vaginocutaneous fistula and inguinal abscess presented 6 years after tension-free vaginal tape sling. *Arch. Ital. Urol. Androl.* **85**, 104–106 (2013).
336. Estevez, J. P., Cosson, M. & Boukerrou, M. An uncommon case of urethrovaginal fistula resulting from tension-free vaginal tape. *Int. Urogynecol. J.* **21**, 889–891 (2010).
337. Karabulut, A., Demitas, O. & Gok, S. A late complication of transobturator tape procedure: vaginocutaneous fistula formation with vaginal mesh erosion. *Int. Urogynecol. J.* **25**, 559–561 (2014).
338. Kobashi, K. C. et al. Erosion of woven polyester pubovaginal sling. *J. Urol.* **162**, 2070–2072 (1999).
339. Yamada, B. S., Govier, F. E., Stefanovic, K. B. & Kobashi, K. C. Vesicovaginal fistula and mesh erosion after Perigee (transobturator polypropylene mesh anterior repair). *Urology* **68**, 1121 e5–e7 (2006).
340. Usher, F., Cogan, J. & Lowry, T. A new technique for the repair of inguinal and incisional hernias. *Arch. Surg.* **81**, 847 (1960).
341. Costello, C., Bachman, S., Grant, S., Cleveland, D. & Loy, T. Characterization of heavyweight and lightweight polypropylene prosthetic mesh explants from a single patient. *Surg. Innov.* **14**, 168–176 (2007).
342. Klinge, U., Klosterhalfen, B., Müller, M. & Schumpelick, V. Foreign body reaction to meshes used for the repair of abdominal wall hernias. *Eur. J. Surg.* **165**, 665–673 (1999).
343. Klosterhalfen, B., Junge, K., Hermanns, B. & Klinge, U. Influence of implantation interval on the long-term biocompatibility of surgical mesh. *Br. J. Surg.* **89**, 1043–1048 (2002).
344. Patel, H., Ostergard, D. R. & Sternschuss G. Polypropylene mesh and the host response. *Int. Urogynecol. J.* **23**, 669–679 (2012).
345. Langhans, T. Ueber riesenzellen mit wandständigen kernen in tuberkeln und die fibröse form des tuberkels [German]. *Virchows Arch. Pathol. Anat.* **42**, 382–404 (1868).
346. Metchnikoff, E. Untersuchungen über die intracelluläre Verdauung wirbellosen bei Tieren [German]. *Arb. Zool. Inst. Univ. Wien* **5**, 141–168 (1884).
347. Tang, L. & Eaton, J. Natural responses to unnatural materials: A molecular mechanism for foreign body reactions. *Mol. Med.* **5**, 351–358 (1999).
348. Tang, L. & Eaton, J. Fibrin(ogen) mediates acute inflammatory responses to biomaterials. *J. Exp. Med.* **178**, 2147–2156 (1993).
349. McNally, A. & Anderson, J. Interleukin-4 induces foreign body giant cells from human monocytes/macrophages. Differential lymphokine regulation of macrophage fusion leads to morphological variants of multinucleated giant cells. *Am. J. Pathol.* **147**, 1487–1499 (1995).
350. Laskin, D. & Pendino, K. Macrophages and inflammatory mediators in tissue injury. *Annu. Rev. Pharmacol. Toxicol.* **35**, 655–77 (1995).
351. Anderson, J. & Miller, K. Biomaterial biocompatibility and the macrophage. *Biomaterials* **5**, 5–10 (1984).
352. Anderson, J. Biological responses to materials. *Annu. Rev. Mater. Res.* **31**, 81–110 (2001).
353. Elmer, C., Blomgren, B., Falconer, C., Zhang, A. & Altman, D. Histological inflammatory response to transvaginal polypropylene mesh for pelvic reconstructive surgery. *J. Urol.* **181**, 1189–1195 (2009).
354. Kavvadias, T., Kaemmer, D., Klinge, U., Kuschel, S. & Schuessler, B. Foreign body reaction in vaginally eroded and noneroded polypropylene suburethral slings in the female: a case series. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **20**, 1473–1476 (2009).
355. Reinke, J. & Sorg, H. Wound repair and regeneration. *Eur. Surg. Res.* **49**, 35–43 (2012).
356. Manodoro, S. et al. Graft-related complications and biaxial tensiometry following experimental vaginal implantation of flat mesh of variable dimensions. *BJOG* **120**, 244–250 (2013).
357. Gonzalez, R. & Ramshaw, B. Comparison of tissue integration between polyester and polypropylene prostheses in the preperitoneal space. *Am. Surg.* **69**, 471–476 (2003).
358. Jongebloed, W. & Worst, J. Degradation of polypropylene in the human eye: a SEM-study. *Doc. Ophthalmol.* **64**, 143–152 (1986).
359. Clave, A. et al. Polypropylene as a reinforcement in pelvic surgery is not inert comparative analysis of 100 explants. *Int. Urogynecol. J.* **21**, 261–270 (2010).
360. Cozad, M. et al. Materials characterization of explanted polypropylene, polyethylene terephthalate and expanded polytetrafluoroethylene composites: spectral and thermal analysis. *J. Biomed. Mater. Res.* **94**, 455–462 (2010).
361. Wood, A. et al. Materials characterization and histological analysis of explanted polypropylene, PTFE, and PET hernia meshes from an individual patient. *J. Mater. Sci. Mater. Med.* **24**, 1113–1122 (2013).
362. Keys, T., Aboushwareb, T. & Badlani, G. Re: post-implantation alterations of polypropylene in the human. *J. Urol.* **189**, 1996–2000 (2013).
363. Rosa, D., Angelini, J., Agnelli, J. & Mei, L. The use of optical microscopy to follow the degradation of isotactic polypropylene (iPP) subjected to natural and accelerated aging. *Polym. Test.* **24**, 1022–1026 (2005).

364. Schmidt, H. *et al.* Comparison of the rates of polypropylene fibre degradation caused by artificial light and sunlight. *Fibres Text. East Eur.* **4**, 53–58 (2011).
365. Blais, P., Carlsson, D., Clark, F., Sturgeon, P. & Wiles, D. The photo-oxidation of polypropylene monofilaments: part II: physical changes and microstructure. *Text. Res. J.* **46**, 641–664 (1976).
366. Liebert, T., Chartoff, R., Cosgrove, S. & McCuskey, R. Subcutaneous implants of polypropylene filaments. *J. Biomed. Mater. Res.* **10**, 939–951 (1976).
367. Bertin, D., Leblanc, M., Marque, S. & Siri, D. Polypropylene degradation: Theoretical and experimental investigations. *Polym. Degrad. Stabil.* **95**, 782–791 (2010).
368. Otto, J. *et al.* Large-pore PDS mesh compared to small-pore PG mesh. *J. Invest. Surg.* **23**, 190–196 (2010).
369. Kidd, B. & Urban, L. Mechanisms of inflammatory pain. *Br. J. Anaesth.* **87**, 3–11 (2001).
370. Miller, J., Acar, F., Kaimaktchiev, V., Gultekin, S. & Burchiel, K. Pathology of ilioinguinal neuropathy produced by mesh entrapment: case report and literature review. *Hernia* **12**, 213–216 (2008).
371. Johner, A., Faulds, J. & Wiseman, S. Planned ilioinguinal nerve excision for prevention of chronic pain after inguinal hernia repair: a meta-analysis. *Surgery* **150**, 534–541 (2011).
372. Kumar, V., Abbas, A., Aster, J. & Fausto, N. *Robbins and Cotran pathologic basis of disease*, 8th edn (Saunders, 2010).
373. Chin, Y. & Stanton, S. A follow up of silastic sling for genuine stress incontinence. *Br. J. Obstet. Gynaecol.* **102**, 143–147 (1995).
374. Duckett, J. & Constantine, G. Complications of silicone sling insertion for stress urinary incontinence. *J. Urol.* **163**, 1835–1837 (2000).
375. Bent, A., Ostergard, D. & Zwick-Zaffuto, M. Tissue reaction to expanded polytetrafluoroethylene suburethral sling for urinary incontinence: clinical and histologic study. *Am. J. Obstet. Gynecol.* **169**, 1198–1204 (1993).
376. Pierce, L. *et al.* Long-term histologic response to synthetic and biologic graft materials implanted in the vagina and abdomen of a rabbit model. *Am. J. Obstet. Gynecol.* **200**, 546.e1–e8 (2009).
377. Hamouda, A., Kennedy, J., Grant, N., Nigam, A. & Karanjia, N. Mesh erosion into the urinary bladder following laparoscopic inguinal hernia repair; is this the tip of the iceberg? *Hernia* **14**, 317–319 (2010).
378. Agrawal, A. & Avill, R. Mesh migration following repair of inguinal hernia: a case report and review of literature. *Hernia* **10**, 79–82 (2006).
379. Elliott, M. & Juler, G. Comparison of Marlex mesh and microporous teflon sheets when used for hernia repair in the experimental animal. *Am. J. Surg.* **137**, 342–344 (1979).
380. Svabik, K. *et al.* Ultrasound appearances after mesh implantation—evidence of mesh contraction or folding? *Int. Urogynecol. J.* **22**, 529–533 (2011).
381. Moalli, P. *et al.* Tensile properties of five commonly used mid-urethral slings relative to the TVT. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* **19**, 655–663 (2008).
382. Welty, G., Klinge, U., Klosterhalfen, B., Kasperk, R. & Schumpelick, V. Functional impairment and complaints following incisional hernia repair with different polypropylene meshes. *Hernia* **5**, 142–147 (2001).
383. Jerabek, J. *et al.* Evaluation of three purely polypropylene meshes of different pore sizes in an onlay position in a New Zealand white rabbit model. *Hernia* **18**, 855–864 (2014).
384. Klinge, U. *et al.* Impact of polymer pore size on the interface scar formation in a rat model. *J. Surg. Res.* **103**, 208–214 (2002).
385. Klinge, U., Otto, J. & Muhl, T. High structural stability of textile implants prevents pore collapse and preserves effective porosity at strain. *Biomed. Res. Int.* **2015**, 953209 (2015).
386. Frostling, H. *et al.* Analytical, occupational and toxicologic aspects of the degradation products of polypropylene plastics. *Scand. J. Work Environ. Health* **10**, 163–169 (1984).
387. Junge, K. *et al.* Persistent extracellular matrix remodelling at the interface to polymers used for hernia repair. *Eur. Surg. Res.* **35**, 497–504 (2003).
388. Birolini, C., Minossi, J., Lima, C., Utiyama, E. & Rasslan, S. Mesh cancer: long-term mesh infection leading to squamous-cell carcinoma of the abdominal wall. *Hernia* **18**, 897–901 (2014).
389. The World Health Organization Internal Agency For Research on Cancer. *IARC Monographs on the evaluation of carcinogenic risks to humans: surgical implants and other foreign bodies* (IARC Press, 1999).
390. Witherspoon, P., Bryson, G., Wright, D., Reid, R. & O'Dwyer, P. Carcinogenic potential of commonly used hernia repair prostheses in an experimental model. *Br. J. Surg.* **91**, 368–372 (2004).
391. Blanchard, K. *et al.* Transponder-induced sarcoma in the heterozygous p53± mouse. *Toxicol. Pathol.* **27**, 519–527 (1999).
392. Carminato, A., Vascellari, M., Marchioro, W., Melchioti, E. & Mutinelli, F. Microchip-associated fibrosarcoma in a cat. *Vet. Dermatol.* **22**, 565–569 (2011).
393. Albrecht, K. Microchip-induced tumors in laboratory rodents and dogs: A review of the literature 1990–2006 in *Technology and Society (ISTAS)*, 2010 IEEE International Symposium on 337–349 (2010).
394. Vascellari, M., Melchioti, E. & Mutinelli, F. Fibrosarcoma with typical features of postinjection sarcoma at site of microchip implant in a dog: histologic and immunohistochemical study. *Vet. Pathol.* **43**, 545–548 (2006).
395. Taylor, C., Siddiqi, I. & Brody, G. Anaplastic large cell lymphoma occurring in association with breast implants: review of pathologic and immunohistochemical features in 103 cases. *Appl. Immunohistochem. Mol. Morphol.* **21**, 13–20 (2013).
396. Thompson, P. & Prince, H. Breast implant-associated anaplastic large cell lymphoma: a systematic review of the literature and mini-meta analysis. *Curr. Haematol. Malign. Rep.* **8**, 196–210 (2013).
397. Sagan, C. *The fine art of baloney detection*, 201–219 (Ballantine Books, 1996).

Author contributions

All authors made a researched data for this article, J.G.B., M.S.B., G.M., R.B. and V.I. contributed to discussions of content, J.G.B., R.S.P., M.S.B., G.M., R.B. and V.I. wrote the manuscript and J.G.B., R.S.P., M.S.B., R.B. and V.I. made a substantial contribution to reviewing and/or editing of this manuscript prior to submission.